

Q9/787360

532 Rec'd PCT/PTO 16 MAR 2001

LAW OFFICES

SUGHRUE, MION, ZINN, MACPEAK & SEAS, PLLC

2100 PENNSYLVANIA AVENUE, N.W.

WASHINGTON, DC 20037-3213

TELEPHONE (202) 293-7060

FACSIMILE (202) 293-7860

www.sughrue.com

March 16, 2001

BOX PCTCommissioner for Patents
Washington, D.C. 20231PCT/JP99/05039
-filed September 16, 1999

Re: Application of Masato HORIE, Keiichi OKUTOMI, Yoshihiro TANIGUCHI, Mikio SUZUKI and
Yutaka OHBUCHI
LY6H GENE
Our Ref: Q63396

Dear Sir:

The following documents and fees are submitted herewith in connection with the above application for the purpose of entering the National stage under 35 U.S.C. § 371 and in accordance with Chapter II of the Patent Cooperation Treaty:

- ☒ an executed Declaration and Power of Attorney.
- ☒ an English translation of the International Application with executed Translator's Declaration.
- ☒ one (1) sheet of drawing.
- ☒ three (3) pages of Sequence Listing and a 3.5" disk containing Sequence Listing.
- ☒ Statement to Support Filing and Submission in Accordance with 37 C.F.R. §§ 1.821-1.825
- ☒ an executed Assignment and PTO 1595 form.
- ☒ International Search Report and a Form PTO-1449 listing the ISR references.

It is assumed that copies of the International Application, the International Preliminary Examination Report, and any Articles 19 and 34 amendments as required by § 371(c) will be supplied directly by the International Bureau, but if further copies are needed, the undersigned can easily provide them upon request.

The Government filing fee is calculated as follows:

Total claims	27	-	20	=	7	x	\$18.00	=	\$126.00
Independent claims	5	-	3	=	2	x	\$80.00	=	\$160.00
Base Fee									\$860.00
Multiple Dependent Claim Fee									\$270.00
TOTAL FILING FEE									\$1416.00
Recordation of Assignment									\$ 40.00
TOTAL FEE									\$1456.00

Checks for the statutory filing fee of \$1416.00 and Assignment recordation fee of \$40.00 are attached. You are also directed and authorized to charge or credit any difference or overpayment to Deposit Account No. 19-4880. The Commissioner is hereby authorized to charge any fees under 37 C.F.R. §§ 1.16, 1.17 and 1.492 which may be required during the entire pendency of the application to Deposit Account No. 19-4880. A duplicate copy of this transmittal letter is attached.

Priority is claimed from September 17, 1998 based on Japanese Application No. 1998-263550.

Respectfully submitted,

Waddell A. Biggart
Waddell A. Biggart
Registration No. 24,861

WAB/amt

09/787360

532 Rec'd PCT/PTO 16 MAR 2001

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of

Masato HORIE et al.

Appln. No.: Not Yet Assigned

Group Art Unit: Not Yet Assigned

Filed: March 16, 2001

Examiner: Not Yet Assigned

CONFIRMATION NUMBER: Not Yet Assigned

For: LY6H GENE

STATEMENT TO SUPPORT FILING AND SUBMISSION IN
ACCORDANCE WITH 37 C.F.R. §§ 1.821-1.825

Assistant Commissioner for Patents
Washington, D.C. 20231
Box SEQUENCE

Sir:

In connection with a Sequence Listing submitted concurrently herewith, the undersigned hereby states that:

1. the content of 3-page Sequence Listing being filed with the present application, and the attached computer readable copy of the Sequence Listing, submitted in accordance with 37 C.F.R. § 1.821(c) and (e), respectively, are the same; and
2. all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true, and further, that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United

09/787360


532 Rec'd PCT/PTO 16 MAR 2001

**STATEMENT TO SUPPORT FILING AND SUBMISSION
IN ACCORDANCE WITH 37 C.F.R. §§ 1.821-1.825**

Q63396

States Code and that such willful false statements may jeopardize the validity of the application or any patent resulting therefrom.

Respectfully submitted,



Drew Hissong
Registration No. 44,765

SUGHRUE, MION, ZINN,
MACPEAK & SEAS, PLLC
2100 Pennsylvania Avenue, N.W.
Washington, D.C. 20037-3213
Telephone: (202) 293-7060
Facsimile: (202) 293-7860

Date: March 16, 2001

1/PRTS

-1-

DESCRIPTION

LY6H GENETECHNICAL FIELD

The invention relates to a gene expressed with high
5 specificity at a high level in the brain, more
particularly a gene coding for a novel protein belonging
to the Ly6 Family (cf. the literature cited below) which
has been utilized in the purification of blood stem cells,
studies on the differentiation of blood cells, activation
10 of immune cells, inhibition of production of active
immune cells, treatment of tumors and the like. The
invention further relates to a novel protein encoded by
said gene and to its specific antibody. In addition, the
invention relates to a therapeutic and prophylactic
15 composition for neurodegenerative disease such as
Alzheimer's disease.

BACKGROUND ART

Proteins of the Ly6 family have a low-molecular
weight GPI-anchored structure and have been identified as
20 a class of cell surface glycoproteins forming a gene
cluster on mouse chromosome 15 [Proc. Natl. Acad. Sci.,
USA., 84, 1638-1643 (1987)].

The Ly6 family is specifically expressed at high
levels in bone marrow cells and lymphoid cells and,
25 therefore, has been utilized as a marker for T-cell

differentiation and hematopoietic stem cells [Immunol. Cell Biol., 73, 277-296 (1995)]. While much remains to be known about its functions in vivo, the finding that its expression is highly modulated in the lymphocytic system suggests that these proteins are playing important roles in the immune system, particularly in the differentiation and function of T cells. It is reported that Ly6c, for instance, mediates the homing of CD8⁺ T cells to the lymph node through integrin-dependent adhesion [Proc. Natl. Acad. Sci., USA., 94, 6898-6903 (1997)].

Furthermore, many GPI-anchored proteins are known to interact with protein kinases [Science, 254, 1016-1019 (1991)]. For example, the interaction of Ly6 with p56lck and p59fyn suggests the likelihood of its involvement in the signal transduction of T cells [Eur. J. Immunol., 23, 825-831 (1993)]. It is also reported that T cells derived from Ly6a-defective mice have been enhanced in the ability to proliferate in response to antigenic stimulation [J. Exp. Med., 186, 705-717 (1997)]. The possibility of its regulating not only the activation of T cells but also that of B cells has also been suggested [J. Immunol., 144, 2197-2204 (1990)].

Furthermore, several GPI-anchored proteins are known to have been expressed and be functioning in both the lymphocytic system and the nervous system [Nature,

379, 826-829 (1996); Curr. Biol., 7, 705-708 (1997)]. In the Ly6 family, Ly6a.2 and Ly6E are reportedly present and functioning in both systems [Proc. Natl. Acad. Sci., USA., 85, 2255-2259 (1996); J. Immunol, 157, 969-973 (1996)].

Elucidation of the physiological roles played by such proteins of the Ly6 family and the genes coding for the proteins and the resulting information are considered to be of use in the field of fundamental scientific research as well as in the pharmaceutical field in connection with the purification of blood stem cells, studies on the differentiation of blood cells, activation of immune cells, inhibition of activation of immune cells, therapy of tumors, and the like.

Recently, in patients with Alzheimer's disease, an excessive cerebral temporal lobe atrophy as compared with age-associated brain atrophy has been reported [Jobst, K. A., et al., Lancet, 343, 829-830 (1994)], suggesting that some gene or genes having a bearing on the cerebral temporal lobe are somehow associated with the onset and progression of Alzheimer's disease. It is logical to assume that should such a gene be identified or characterized, there might be provided information useful for the therapy and prophylaxis of Alzheimer's disease.

Therefore, an object of the present invention is to

provide the above information needed by those concerned, particularly a novel human protein belonging to the Ly6 family and a gene coding for the protein.

5 A further object of the invention is to provide a pharmaceutical composition for the therapy and prophylaxis of various neurodegenerative diseases, represented by Alzheimer's disease.

10 The present inventor explored into the genes derived from various human tissues and succeeded in isolating and characterizing a novel brain-specific gene meeting the above objects. The inventor further found that the level of expression of this newly isolated gene is markedly depressed in the temporal lobe, inclusive of the hippocampus and entorhinal cortex, of a patient with
15 Alzheimer's disease, that this is a causative factor in the onset and progression of Alzheimer's disease and in dementia and other disturbances and that this gene and its expression product can be exploited with advantage in the therapy and prophylaxis of Alzheimer's disease. The
20 present invention has been accomplished on the basis of the above findings.

DISCLOSURE OF INVENTION

The present invention provides a gene comprising a nucleotide sequence coding for the following protein (a)
25 or (b).

(a) a protein having the amino acid sequence shown in SEQ ID NO:1

(b) a protein having an amino acid sequence derived from the amino acid sequence shown in SEQ ID NO:1 by deletion, substitution or addition of one or a plurality of amino acids and having at least one physiological activity selected from the group consisting of neuronal survival-supporting activity, nerve elongating activity, nerve regenerating activity, neuroglia-activating activity, and mnemonic (brain memory-forming) activity.

The invention also provides the above gene wherein the nucleotide sequence is shown in SEQ ID NO:2, in particular, which is a human gene.

Furthermore, the invention provides a gene comprising the following polynucleotides (a) or (b), particularly the corresponding human gene.

(a) a polynucleotide containing the nucleotide sequence shown in SEQ ID NO:3

(b) a polynucleotide which hybridizes under stringent condition with a DNA having the nucleotide sequence shown in SEQ ID NO:3.

The invention further provides a gene expression vector harboring said gene; a host cell harboring said gene expression vector; an expression product which is expressed by said host cell; a protein encoded by the

gene of the invention; and an antibody which binds said expression product or said protein.

The invention further provides a therapeutic and prophylactic composition for neurodegenerative disease, which comprises said protein or an equivalent thereof or said expression product as an active ingredient in combination with a pharmaceutical carrier. More particularly, the invention provides the therapeutic and prophylactic composition for neurodegenerative disease, wherein said active ingredient is a protein having the amino acid sequence shown in SEQ ID NO:1 or an equivalent thereof or a gene product obtainable by expression of the whole or part of a gene comprising a nucleotide sequence shown in SEQ ID NO:2 and having at least one physiological action selected from the group consisting of neuronal survival-supporting action, nerve elongating action, nerve-regenerating action, neuroglia-activating action, and brain memory-forming (mnemonic, encoding) action.

Especially, the invention provides the therapeutic and prophylactic composition for Alzheimer's disease, Alzheimer type dementia, brain ischemia and Parkinson's disease.

In addition, the invention provides a sense strand oligonucleotide comprising at least 20 consecutive

constituent nucleotides of the nucleotide sequence shown
in SEQ ID NO:2; a gene therapy composition comprising
said sense strand oligonucleotide as an active ingredient
in combination with a pharmaceutical carrier; and a gene-
5 specific probe comprising an oligonucleotide sequence of
at least 10 consecutive constituent nucleotides of the
nucleotide sequence shown in SEQ ID NO:2.

Furthermore, the invention provides a method of
screening for candidate compounds either capable of
10 binding to said protein, equivalent thereof or expression
product or influencing its activity which comprises using
said protein, equivalent or expression product; a kit for
said screening; and said compounds so screened.

Representation of amino acids, peptides, nucleotide
15 sequences, nucleotides, etc. by abbreviations in the
specification is in conformity with the rules recommended
by IUPAC-IUB [IUPAC-IUB Communication on Biological
Nomenclature, Eur. J. Biochem., 138, 9 (1984)],
"Guideline for drafting patent specifications relative to
20 nucleotide sequences and/or amino acid sequences" (edited
by the Patent Office of Japan) and the conventions
relating to the use of codes or symbols in the art.

A specific example of the gene of the invention is
the gene deduced from the DNA sequence of the PCR product
25 designated "LY6H" as described in the example which

appears later herein. Its nucleotide sequence is as shown in SEQ ID NO:3.

The LY6H gene is a cDNA containing a 420-codon open reading frame (ORF) coding for a novel brain-specific
5 protein (LY6H protein) having a 140-residue amino acid sequence as shown in SEQ ID NO:1, and having a full-length sequence of 854 nucleotides.

The LY6H protein which is the expression product of the gene of the invention was found to have high homology
10 to mouse Ly6 family proteins [Immunol. Cell Biol., 73, 277-296 (1995)] by a GenBank/EMBL database search using FASTA Program (Person, W. R. et al., Proc. Natl. Acad. Sci., USA., 85, 2444-2448 (1988)). Furthermore, high gene-to-gene homology was recognized. Therefore, the gene
15 of the invention is considered to be a novel human Ly6 gene.

The LY6 gene of the invention was identified to be a gene which is specifically expressed in the brain by the sequencing of more than 28000 cDNA clones selected at
20 random from a fetal human brain cDNA library. By RH chromosome mapping [Hum. Mol. Genet., 5, 339-346 (1996)], the locus of the gene on the chromosome was found to be 8q24. 3.

Thus, the gene and expression product of the
25 invention, thus provided, contribute to detection of the

expression of the gene in various tissues, production of human LY6H protein by genetic engineering techniques, and construction of an antibody thereto, hence enabling the purification of hematopoietic stem cells, study of blood
5 cell differentiation, activation or suppression of immune cells, therapy of tumors, and the like.

In addition, the expression product (polypeptide) of the invention, thus provided, enables provision of a drug for prophylaxis and therapy of neurodegenerative
10 diseases such as Alzheimer's disease, Alzheimer type dementia, Parkinson's disease and ischemic brain. Furthermore, the sense strand of the gene according to the invention can be utilized as a pharmaceutical composition for gene therapy, with which the onset and
15 progression of the above-mentioned neurodegenerative diseases can be inhibited or arrested.

The invention further provides a method of screening for compounds either binding or influencing the activity of the expression product (polypeptide) of the
20 invention and a relevant kit for screening, hence compounds so screened as well. For identification of such compounds screened, an antibody binding to the expression product of the gene of the invention can be utilized.

In the specification, the term "gene" is used to
25 mean a double-stranded DNA and its constituent single-

stranded DNA, whether sense or antisense, without regard to its length. Therefore, unless otherwise indicated, the gene (DNA) of the invention includes a double-stranded DNA containing a human genomic DNA, a single-stranded DNA (sense strand) inclusive of the cDNA, a single-stranded DNA (antisense strand) having a sequence complementary to said sense strand, and fragments of said DNAs.

The gene (DNA) of the invention may contain a leader sequence, a coding region, exons and introns. The polynucleotide includes both RNA and DNA. The DNA includes cDNA, genomic DNA and synthetic DNA. The polypeptide includes its fragments, homologs, derivatives and mutants. The mutants include alleles which occur naturally, mutants not existing naturally, mutants having amino acid sequences mutated by deletion, substitution, addition and/or insertion, and mutants having functionally equivalent modified amino acid sequences.

Such modifications (e.g. mutations) of amino acid sequences may for example occur from spontaneous mutation or posttranslational modification but may be artificially induced by utilizing a native gene (for example, specific genes of the invention).

The homology of such mutants to the unmutated polypeptide may be at least 70%, preferably 80%, more preferably 95%, still more preferably 97%. The above

polypeptide and its mutants and homologs have a structural feature conserved in common and may have the biological activities of the expression product of the gene of the invention, such as neuronal survival-

5 supporting action, neuronal growth-stimulating action, nerve generating action, and neuroglia-stimulating action. The homology of polypeptides can be analyzed by searching through a database such as SWISSPLOTS Database using a sequence analysis software such as FASTA [Clustal, V.,
10 Methods Mol. Biol., 25, 307-318 (1994)].

The gene coding for such a mutant is silent or conserved for amino acid substitution. Thus, the amino acid residues encoded by the nucleotide sequence are not altered.

15 The conservatively substitutable amino acid residues, i.e. the amino acid residues substitutable with other amino acid residues without losing the activities of the polypeptide having such original amino acid residues, and the corresponding original amino acid
20 residues are as follows.

Original amino acid residue	Conservatively substituting amino acid residue
Ala	Ser
25 Arg	Lys
Asn	Gln, His

	Asp	Glu
	Cys	Ser
	Gln	Asn
	Glu	Asp
5	Gly	Pro
	His	Asn or Gln
	Ile	Leu or Val
	Leu	Ile or Val
	Lys	Arg, Aln or Glu
10	Met	Leu or Ile
	Phe	Met, Leu or Tyr
	Ser	Thr
	Thr	Ser
	Trp	Tyr
15	Tyr	Trp or Phe
	Val	Ile or Leu

In addition, Cys may be substituted for a different kind of amino acid residue, e.g. Ser, Ala or Val.

The gene and expression product according to the invention provide information and means of great use for the elucidation, expatiation, diagnosis, prophylaxis and therapy of neurodegenerative diseases such as Alzheimer's disease, brain ischemia and Parkinson's disease. The gene of the invention can also be used with advantage for the development of new drugs capable of inducing expression

of the gene for use in the treatment of said neurode-
generative diseases. In addition, the detection of
expression of the gene of the invention and of the
resulting expression product in individuals or tissues,
5 and the detection of mutation (deletion or point
mutation) of the gene or abnormal expression thereof can
be utilized with advantage for the elucidation and
diagnosis of said neurodegenerative diseases.

The gene of the invention includes but is not
10 limited to the gene having the nucleotide sequence shown
in SEQ ID NO:2 which codes for a protein having the amino
acid sequence shown in SEQ ID NO:1, for example a gene
(LY6H gene) having the nucleotide sequence shown in SEQ
ID NO:3. For example, the gene of the invention may be a
15 gene coding for an amino acid sequence derived from the
above-defined amino acid sequence by a given modification,
a gene coding for an amino acid sequence having a given
degree of homology to the above-defined amino acid
sequence, or a gene having a nucleotide sequence having a
20 given degree of homology to any of the above genes.

The above-mentioned given degree of homology to a
defined amino acid sequence or nucleotide sequence means
a homology of at least not less than 70%, preferably not
less than 90%, more preferably not less than 95%, still
25 more preferably not less than 97%. The present invention

encompasses homologs (gene homologs and protein homologs) having such homology.

The gene of the invention includes "a gene coding for a polypeptide having an amino acid sequence (modified amino acid sequence) derived from the amino acid sequence shown in SEQ ID NO:1 by deletion, substitution or addition of one or a plurality of amino acids". The extent and position or positions of "deletion, substitution or addition" are not particularly restricted insofar as the resulting polypeptide having a modified amino acid sequence is equivalent, in biological function, to the polypeptide (LY6H protein) having the amino acid sequence shown in SEQ ID NO:1. The biological "function" mentioned above includes physiological functions such as neuronal survival-supporting action, nerve elongating action, nerve regenerating action, neuroglia-activating action, and mnemonic action, and the "equivalent" is a polypeptide having such functions. Therefore, the protein having such a modified amino acid sequence includes a protein (equivalent) having a fragment (a consecutive-residue fraction) of the amino acid sequence shown in SEQ ID NO:1 and having physiological activities similar to those mentioned above for the full-length of said amino acid sequence. Furthermore, the gene coding for a polypeptide having the above-modified amino acid sequence

may be a gene with which the gene of the invention
encoding a polypeptide having the pre-modification amino
acid sequence can be detected. The term plurality as used
in connection with said modification usually means not
5 less than 2 but up to several, although the range is not
restrictive.

The homolog of LY6H gene (and the homolog of the
expression product of the gene) according to the
invention means any of a series of related genes (and
10 their expression products) which are homologous in
sequence and recognized as one gene family from their
structural characteristics, common gene expression
pattern, and similarities in said biological function.
This, of course, includes alleles of the genes of the
15 invention.

The modification (mutation) of an amino acid
sequence may occur naturally, for example by spontaneous
mutation and posttranslational modification but may be
induced artificially on the basis of the native gene (for
20 example the specific gene of the invention). The
invention covers any and all modified genes having the
above characteristics without regard to the cause or
means of modification or mutation.

The artificial means for said modification
25 (mutation) of the amino acid sequence includes genetic

engineering techniques such as site-specific mutagenesis
[Methods in Enzymology, 154, 350, 367-382 (1987); *ibid.*,
100, 468 (1983); Nucleic Acids Res., 12, 9441 (1984);
"Zoku Seikagaku Jikken Koza (Experiments in Biochemistry,
5 Second Series) 1": Idenshi Kenkyuho (Methods in Gene
Research) II, the Biochemical Society of Japan (ed.), p
105 (1986), etc.], methods of chemical synthesis such as
the phosphotriester method and phosphoamidite method [J.
Am. Chem. Soc., 89, 4801 (1967); *ibid.*, 91, 3350 (1969);
10 Science, 150, 178 (1968); Tetrahedron Lett., 22, 1859
(1981); *ibid.*, 24, 245 (1983)], and combinations of such
methods.

More particularly, the DNA can be synthesized by a
chemical method such as the phosphoamidite method or the
15 phosphotriester method, and this synthesis can be
effected on a commercially available automated oligo-
nucleotide synthesizer. The double-stranded fragment can
be obtained from the chemically synthesized single-strand
fragment by synthesizing a complementary strand and
20 annealing them under suitable conditions or adding the
complementary strand using a suitable primer sequence and
a DNA polymerase.

A specific example of the gene according to the
invention is the gene having the nucleotide sequence
25 shown in SEQ ID NO:3. The coding region (the sequence

shown in SEQ ID NO:2) of this nucleotide sequence is an example of combination of codons specifying the respective amino acid residues of the amino acid sequence shown under SEQ ID NO:1. The gene of the invention is not
5 limited to the gene having said defined nucleotide sequence but includes any gene having a nucleotide sequence obtainable by selecting any arbitrary combination of codons for each amino acid residue. Selection of codons can be made in the routine manner,
10 with reference to the codon usage in the host to be employed [Nucleic Acids Res., 9, 43 (1981)].

Furthermore, the gene of the invention includes one having a nucleotide sequence showing a certain level of homology to the nucleotide sequence shown in SEQ ID NO:3.
15 Inferred by said level of homology are polynucleotides and complementary polynucleotides having at least 70% homology, preferably at least 90% homology, more preferably at least 95% homology, to the nucleotide sequence shown in SEQ ID NO:3. The gene having such a
20 level of homology may for example be characterized as a polynucleotide which hybridizes with a DNA having the nucleotide sequence shown in SEQ ID NO:3 under stringent conditions. More particularly, the gene having a nucleotide sequence which hybridizes with the DNA having
25 the nucleotide sequence shown in SEQ ID NO:3 under the

condition of 6 × SSC at 65°C overnight or 50% formamide-4
× SSC at 37°C overnight is subsumed in the concept of the
gene having said level of homology. Here, SSC stands for
standard saline citrate (1 × SSC = 0.15 M NaCl, 0.015 M
5 sodium citrate).

The gene of the invention can be easily produced
and isolated by the general genetic engineering
technology based on the sequence information on any
specific example of the gene of the invention as
10 disclosed in this specification [e.g. Molecular Cloning
2d Ed, Cold Spring Harbor Lab. Press (1989); Zoku
Seikagaku Jikken Koza (Experiments in Biochemistry,
Second Series): "Idenshi Kenkyuho (Methods in Gene
Research) I, II, III, the Biochemical Society of Japan
15 (ed.), (1986)].

More particularly, this can be done by preparing a
cDNA library from a suitable source, in which the gene of
the invention can be expressed, by a routine procedure
and selecting a desired clone from this library using a
20 suitable probe or antibody specific to the gene of the
invention [Proc. Natl. Acad. Sci., USA., 78, 6613 (1981);
Science, 222, 778 (1983)].

The source of cDNA, which can be used in the above
procedure includes various cells and tissues expressing
25 the gene of the invention, as well as cultured cells

derived therefrom, particularly brain tissues. Isolation of the total RNA from such a source, isolation and purification of mRNA, and acquisition and cloning of cDNA can also be carried out in the conventional manner.

- 5 Moreover, cDNA libraries are commercially available and the present invention can be carried into practice using such cDNA libraries, for example those cDNA libraries available from CLONTECH Lab. Inc.

10 The method of screening for the gene of the invention from a cDNA library is not particularly restricted but the conventional procedure can be employed. Examples of the screening methods include an immuno-screening method using a specific antibody to the protein produced by a cDNA to select the corresponding cDNA clone,
15 a method using a probe selectively binding to the objective DNA sequence, such as a plaque hybridization method, and a colony hybridization method, and a combination of such methods.

As the probe for the above method, the DNA
20 chemically synthesized according to the nucleotide sequence information on the gene of the invention can be generally employed. The gene of the invention which has already been obtained or a fragment thereof can also be used as the probe with advantage. The sense primer and
25 antisense primer established according to the nucleotide

sequence information on the gene of the invention can be used as screening probes.

The nucleotide sequence for use as the probe may be a partial nucleotide sequence corresponding to SEQ ID NO:2 and comprising at least 10 consecutive nucleotides, preferably 20 consecutive nucleotides, more preferably 30 consecutive nucleotides, most preferably 50 consecutive nucleotides. Moreover, the positive clone having the oligonucleotide sequence shown in SEQ ID NO:2 as such can be used as the probe.

In obtaining the gene of the invention, the DNA/RNA amplification by PCR [Science, 230, 1350 (1985)] can be used with advantage. Particularly when a full-length cDNA can hardly be obtained from a library, the RACE method [Rapid amplification of cDNA ends; Jikken Igaku (Experimental Medicine), 12(6), 35 (1994)], especially 5'-RACE method [M. A. Frohman, et al., Proc. Natl. Acad. Sci., USA., 8, 8998 (1988)], can be used with advantage.

The primers for use in such PCR methods can be judiciously established with reference to the sequence information on the gene of the invention as disclosed herein and can be synthesized by the routine procedure. The isolation and purification of the amplified DNA/RNA fragment can be carried out in the routine manner as mentioned above, for example by the gel electrophoresis

method.

Sequencing of the gene of the invention as obtained in the above manner or various DNA fragments can be made in accordance with the dideoxy method [Proc. Natl. Acad. Sci., USA., 74, 5463 (1977)] or the Maxam and Gilbert method [Methods in Enzymology, 65, 499 (1980)] or more expediently by using a commercial sequencing kit.

For example, with the gene of the invention thus obtained, the expression or non-expression of the gene of the invention in an individual or a given tissue can be specifically detected by utilizing a portion or the whole of the nucleotide sequence of the gene of the invention.

The above detection can be made by the conventional procedures, such as RNA amplification by RT-PCR [reverse transcribed-polymerase chain reaction; E.S. Kawasaki, et al., Amplification of RNA. In PCR Protocol. A Guide to Methods and Applications, Academic Press, Inc., San Diego, 21-27 (1991)], Northern blot analysis [Molecular Cloning, Cold Spring Harbor Lab. (1989)], determination on cellular level by in situ RT-PCR [Nucl. Acids Res., 21, 3159-3166 (1993)] or in situ hybridization, NASBA [nucleic acid sequence-based amplification, Nature, 350, 91-92 (1991)], and the like conventional techniques. The preferred is the RT-PCR detection method.

The primer which is to be used when the PCR method

is chosen for the above purpose is not particularly restricted insofar as it is characteristic of the gene of the invention and capable of selective amplification of the particular gene only and can be judiciously

5 established based on the sequence information on the gene of the invention. Usually, one having a partial sequence of the gene of the invention, which is about 10-35 nucleotides long, preferably about 15-30 nucleotides long can be used as the primer.

10 The gene of the invention, thus, includes the DNA fragment which can be used as a specific primer and/or specific probe for the detection of the LY6H gene of the invention.

The DNA fragment mentioned above can be defined as
15 a polynucleotide which hybridizes with the polynucleotide having the nucleotide sequence shown in SEQ ID NO:2 under stringent conditions. The stringent conditions mentioned above may be the ordinary conditions for primers or probes and, as such, are not particularly restricted. For
20 example, the above-mentioned conditions of 6 × SSC, 65°C, overnight or the condition of 50% formamide-4 × SSC, 37°C, overnight can be mentioned.

By applying the gene of the invention to the standard genetic engineering technology, the expression
25 product (polypeptide) of the gene or a protein containing

it can be easily produced in large quantities and with good reproducibility.

Therefore, the invention further provides a polypeptide having the amino acid sequence encoded by the gene of the invention (the expression product of the invention), a vector harboring the gene of the invention for the production of the polypeptide, a host cell transfected with the vector, and a method of producing the polypeptide of the invention which comprises growing the host cell.

The polypeptide (LY6H protein) having the amino acid sequence shown under SEQ ID NO:1 is a specific embodiment of the polypeptide of the invention. The polypeptide of the invention is not limited to this LY6H protein but includes its homolog. The homolog may be a polypeptide having an amino acid sequence derived from the amino acid sequence shown in SEQ ID NO:1 by the deletion, substitution or addition of one or more amino acids and retaining the same function as the LY6H protein. A specific example of the homolog is the expression product of a homolog of said LY6H gene (the LY6H equivalent gene inclusive of the allele).

Furthermore, the homolog of the LY6H protein of the invention includes proteins having the same activity or function as the polypeptide having the amino acid

sequence shown in SEQ ID NO:1 as derived from any of mammals such as equine, sheep, bovine, canine, monkey, cat, bear, etc. and rodents such as rat, mouse and rabbit.

The polypeptide of the invention can be produced by
5 the conventional recombinant DNA technology [e.g. Science, 224, 1431 (1984); Biochem. Biophys. Res. Comm., 130, 692 (1985); Proc. Natl. Acad. Sci., USA., 80, 5990 (1983)] based on the gene sequence information provided by the present invention.

10 More particularly, the production of said polypeptide is carried out by the procedure comprising constructing a recombinant DNA (expression vector) which permits expression of the gene coding for the desired protein in a host cell, transforming the host cell with
15 the vector, growing the resulting transformant, and harvesting the polypeptide from the culture broth.

The host cell mentioned above may be whichever of a prokaryotic cell and an eukaryotic cell. As the prokaryotic host, Escherichia coli, Bacillus subtilis
20 and other common bacteria can be mentioned and preferably cells of Escherichia coli, particularly cells of Escherichia coli K12, can be employed. The eukaryotic host cell includes cells of vertebrates and yeasts and the former include the monkey cell line COS [Cell, 23:
25 175 (1981)], Chinese hamster ovarian cells, and the

dihydrofolate reductase-defective cells thereof [Proc. Natl. Acad. Sci., USA., 77: 4216 (1980)]. As the latter, yeast cells of the genus Saccharomyces can be used with advantage, but these are not exclusive choices.

5 When prokaryotic cells are used as host cells, an expression plasmid construct prepared by using a vector which is replicatable in the particular host cell and adding a promoter and SD (Shine and Dalgarno) sequence upstream of the gene of the invention so that the gene
10 may be expressed therein as well as an initiation codon (e.g. ATG) necessary for initiation of protein synthesis can be used with advantage. As the vector mentioned above, it is usual to employ plasmids derived from Escherichia coli, such as pBR322, pBR325, pUC12, pUC13, etc. However,
15 these are not exclusive choices but various known vectors can be utilized. Examples of the commercial vectors for use in expression systems using E. coli include pGEX-4T (Amersham Pharmacia Biotech), pMAL-C2, pMA1-P2 (New England Biolabs), pET21, pET21/lacq (Invitrogen) and
20 pBAD/His (Invitrogen).

 As the expression vector for use when cells of a vertebrate are used as host cells, the vector having a promoter upstream of the gene of the invention to be expressed, RNA splice sites, polyadenylation site and a
25 transcription termination sequence is usually employed,

and this vector may further have a replication origin where necessary. A specific example of the expression vector is pSV2dhfr harboring an early promoter of SV40 [Mol. Cell. Biol., 1: 854 (1981)]. Aside from the above, various known vectors available commercially can be employed. Examples of the commercial vectors which are used in expression systems using animal cells include vectors for animal cells, such as pEGFP-N, pEGFP-C (CLONTECH), pIND (Invitrogen), pCDNA3.1/His (Invitrogen), etc., and vectors for insect cells, such as pFastBac HT (Gibco BRL), pAcGHLT (PharMingen), pAc5/V5-His, pMT/V5-His and pMT/Bip/V5-his (all Invitrogen).

pAM82 having a promoter for the acid phosphatase gene [Proc. Natl. Acad. Sci., USA., 80: 1 (1983)] is a specific example of the expression vector for use when yeast cells are used as host cells. The commercial expression vectors for yeast cells include pPICZ (Invitrogen) and pPICZ α (Invitrogen).

The promoter is not particularly restricted, either. When a strain of the genus Escherichia is used as the host, tryptophan (trp) promoter, lpp promoter, lac promoter, recA promoter, PL/PR promoter, etc. can be utilized with advantage. When the host is a strain of the genus Bacillus, SP01 promoter, SP02 promoter, penP promoter, etc. are preferably used. When a yeast is used

as the host, pH05 promoter, PGK promoter, GAP promoter, ADH promoter, etc. can be utilized with advantage. The preferred promoter for use when host cells are animal cells include SV40-derived promoters, retrovirus
5 promoters, metallothionein promoter, heat shock promoter, cytomegalovirus promoter, and SR α promoter.

As the expression vector for the gene of the invention, the conventional fusion protein expression vector can be used with advantage. pGEX (Promega) for the
10 expression of glutathione-S-transferase (GST)-fused proteins is a specific example of the vector.

The polynucleotide sequence wherein the coding sequence for a mature polypeptide assists in the expression and secretion of a polypeptide from host cells
15 includes the secretory sequence, the leader sequence and the marker sequence (hexahistidine tag, histidin tag) used in the purification of a fusion mature polypeptide in the case of bacterial cells, and the hemagglutinin (HA) tag in the case of mammalian cells.

20 The method of introducing the recombinant DNA (expression vector) into the host cell and the associated transforming method are not particularly restricted but various standardized methods can be utilized.

The transformant obtained can be cultured in the
25 routine manner, whereby the objective protein encoded by

the deliberately designed gene according to the invention is expressed and produced (accumulated/secreted) intracellularly, extracellularly or on the cell membrane.

The culture medium to be used can be judiciously
5 selected from among various routine media according to the kind of adopted host cell and the culture is also performed under conditions favoring growth of the host cell.

The resulting recombinant protein (LY6H protein)
10 according to the invention can be optionally isolated and purified by various separation techniques taking advantage of its physical and/or chemical properties, for instance ["Seikagaku Data Book (Biochemical Data Book) II", 1175-1259, First Edition, 1st impression, June 23,
15 1980, Tokyo Kagaku Dojin K.K.; Biochemistry, 25(25), 8274 (1986); Eur. J. Biochem., 163, 313 (1987), etc.].

Examples of such techniques are the conventional reconstitution method, treatment with a protein precipitating agent (salting-out method), centrifugation,
20 osmotic shock method, sonic disruption, ultrafiltration, various types of chromatography such as molecular sieve chromatography (gel filtration), adsorption chromatography, ion exchange chromatography, affinity chromatography and high performance liquid chromatography
25 (HPLC), dialysis, and combinations of these techniques.

The particularly preferred technique includes affinity chromatography using a column to which a specific antibody to the protein of the invention has been coupled.

In designing the objective gene encoding the polypeptide of the invention, the nucleotide sequence of Y6H gene as shown in SEQ ID NO:2 can be utilized with advantage. If desired, this gene can be used after the codons specifying the respective amino acid residues have been judiciously altered. Furthermore, when any amino acid residue or partial sequence of the amino acid sequence encoded by the LY6H gene is to be modified by substitution, deletion or addition, such modifications can be made by the various methods described above, for example by site-specific mutagenesis.

The polypeptide of the invention can be produced by the standard protocol for chemical synthesis according to the amino acid sequence shown in SEQ ID NO:1. The method includes the conventional liquid-phase method and solid-phase method for peptide synthesis.

More particularly, the method for peptide synthesis includes the so-called stepwise elongation method in which the constituent amino acids are coupled one by one for chain extension and the fragment condensation method which comprises synthesizing fragments each consisting of several amino acids beforehand and coupling the fragments

together. The synthesis of the protein of the invention can be carried out by whichever of the above two methods.

The method of condensation for use in the above peptide synthesis may also be a conventional one, including the azide process, mixed acid anhydride process, DCC process, active ester process, redox process, DPPA (diphenylphosphoryl azide) process, DCC + additive (1-hydroxybenzotriazole, N-hydroxysuccinamide, N-hydroxy-5-norbornene-2,3-dicarboximide or the like) process and Woodward's reagent process.

The solvent to be used in these processes can also be judiciously selected from among the common solvents well known in the art for use in such peptide-forming condensation reactions. Examples of the solvents include dimethylformamide (DMF), dimethyl sulfoxide (DMSO), hexaphosphoramide, dioxane, tetrahydrofuran (THF), ethyl acetate, etc., and mixtures thereof.

In conducting the peptide synthesizing reactions, the carboxyl group of any amino acid or fragment peptide that should not take part in the reaction can be protected in advance, generally by esterification in the form of a lower alkyl ester such as methyl ester, ethyl ester, tert-butyl ester, etc. or an aralkyl ester such as benzyl ester, p-methoxybenzyl ester, p-nitrobenzyl ester, etc.

Referring to any amino acid having a functional group in its side chain, the hydroxyl group of a tyrosine residue, for instance, may be protected in advance with an acetyl, benzyl, benzyloxycarbonyl, tertiary butyl or
5 other group, although such protection is not necessarily indispensable. Furthermore, the guanidino group of an arginine residue can be protected with a suitable protective group such as nitro, tosyl, p-methoxybenzene-sulfonyl, methylene-2-sulfonyl, benzyloxycarbonyl,
10 isobornyloxycarbonyl, adamantyloxycarboxyl or the like.

The reactions for eliminating such protective groups from the protected amino acids, peptides or the end product protein of the invention can also be carried out in the routine manner, for example by catalytic
15 reduction or a method using liquid ammonia/sodium, hydrogen fluoride, hydrogen bromide, hydrogen chloride, trifluoroacetic acid, acetic acid, formic acid, methane-sulfonic acid or other reagent.

The polypeptide of the invention, thus produced,
20 can be purified as needed by the various techniques mentioned above, such as ion exchange resin chromatography, partition chromatography, gel chromatography, countercurrent distribution and the like methods in routine use in the field of peptide chemistry.

25 The polypeptide of the invention can be used with

advantage as an immunogen for preparation of its specific antibody. By utilizing this immunogen, the antiserum (polyclonal antibody) and the monoclonal antibody can be provided.

5 The technology of producing antibodies is well known to those skilled in the art and the known procedures can be employed in the present invention [e.g. Zoku Seikagaku Jikken Koza (Experiments in Biochemistry, second series) "Men-eki Seikagaku Kenkyuho (Methods in
10 Immunobiochemistry)", edited by the Biochemical Society of Japan (1986)].

For example, as the immune animal for harvesting the desired antiserum therefrom, the ordinary animals such as rabbit, guinea pig, rat, mouse, chicken, etc. can
15 be arbitrarily selected and the immunization with said immunogen and the collection of blood can also be carried out by the conventional procedures.

Preparation of a monoclonal antibody can also be carried out by the conventional technique which comprises
20 constructing a hybridoma between the plasma cell (immune cell) of an animal immunized with said immunogen and a plasmacytoma cell, selecting clones producing the desired antibody, and cultivating the clones. The immune animal is generally selected in consideration of its
25 compatibility with the plasmacytoma cell to be used for

cell fusion and usually the mouse or the rat is used with advantage. The immunization procedure may be the same as used for the preparation of said antiserum and, if desired, the immunization can be made using a
5 conventional adjuvant in combination.

The plasmacytoma cell for use in said hybridization is not particularly restricted, either, but includes various myeloma cells such as p3 (p3/x63-Ag8) [Nature, 256: 495-497 (1975)], p3-U1 [Current Topics in
10 Microbiology and Immunology, 81: 1-7 (1978)], NS-1 [Eur. J. Immunol., 6: 511-519 (1976)], MPC-11 [Cell, 8: 405-415 (1976)], SP2/0 [Nature, 276: 269-271 (1978)], etc., R210 [Nature, 277: 131-133 (1979)] and others in rats, and cells derived therefrom.

15 The hybridization between said immune cell and said plasmacytoma cell can be effected by the known technology in the presence of a conventional hybridization promoter such as polyethylene glycol (PEG) or Sendai virus (HVJ) and the separation of the objective hybridoma can also be
20 carried out in the known manner [Meth. in Enzymol., 73: 3 (1981); Zoku Seikagaku Jikken Koza (ditto)].

The search for the objective antibody-producing cell clone and the monoclonal antibody preparation can also be carried out in the routine manner. For example,
25 the search for the antibody-producing hybridoma can be

made by any of the various techniques in routine use for the detection of antibodies, such as ELISA [Meth. in Enzymol., 70: 419-439 (1980)], plaque method, spot method, agglutination reaction method, Ouchterlony method,
5 radioimmunoassay, and the like, using the protein of the invention as an antigen.

Harvesting of the antibody of the invention from the resulting hybridoma can be achieved by cultivating the hybridoma in the routine manner and recovering the
10 antibody as a culture supernatant or administering the hybridoma to a compatible mammal and recovering the antibody in the form of ascites. The former method is suitable for production of the antibody of high purity, while the latter method is suitable for high-production
15 of the antibody. The antibody thus produced can be further purified by the conventional means such as salting-out, gel filtration, affinity chromatography and the like.

The antibody thus obtained is characterized by its
20 binding affinity for the LY6H protein of the invention and can be used with advantage for the purification of LY6H protein and determination or differentiation of the protein by immunological techniques. Furthermore, since a decreased expression of the gene of the invention has
25 been confirmed in the temporal lobe of the brain of a

patient with Alzheimer's disease which is a neurodegenerative disease, this antibody can be utilized in the screening for agonists or antagonists of LY6H protein.

5 The present invention provides the novel antibody described above, too.

 The polypeptide of the invention is useful in the field of medicine as pharmaceutical products containing it as an active ingredient. Therefore, the invention
10 provides a pharmaceutical composition comprising the polypeptide of the invention as an active ingredient.

 The usefulness of the polypeptide of the invention in or as said pharmaceutical composition is ascribable to the neuronal survival-supporting action, nerve elongating
15 action, nerve regenerating action, neuroglia-activating action and mnemonic action inherent in this brain-specific polypeptide. Examples of the methods for confirming these actions include the following methods for each action.

20 1) Neuronal survival-supporting action

 The following method can be used for quantitating the neuronal survival-supporting action of the polypeptide of the invention. For example, the hippocampus is aseptically isolated from the whole brain
25 of a fetal SD rat and treated with an enzyme, and seeded

in a poly-L-lysine (Sigma)-precoated 96-well plate containing 10% fetal calf serum-DMEM at a final concentration of 2×10^5 cells/cm².

The cells are grown for 24 hours, at the end of which time the culture medium is changed to 1% N2 Supplement (Gibco)-containing DMEM. Then, the active ingredient polypeptide of the invention is added (the invention group). As control, the polypeptide of the invention which has been heat-treated in a boiling water bath for 5 minutes is added (the boiled protein group).

The cells (culture) in each group as prepared in the above manner are cultured for 72 hours. Then, by performing an MTT [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide] assay using Promega Cell Titer 96-Well Assay System, the neuronal survival-supporting effect of the polypeptide of the invention on hippocampal neurons can be evaluated.

Similarly, by isolating the ventral midbrain aseptically from the same whole brain of a fetal SD rat as above and carrying out an MTT assay in the same manner as above, the neuronal survival-supporting effect of the polypeptide of the invention on the midbrain neurons can be investigated.

2) Dopaminergic neuron survival-supporting action

As a method of evaluating the neuronal survival-

supporting action of the polypeptide of the invention,
the following method of quantitating the dopaminergic
neuron survival-supporting activity can be mentioned.
Thus, the cells (culture) in each group as prepared above
5 under 1) are cultured for 72 hours and, then, fixed with
4% paraformaldehyde-PBS by 15-minute standing at room
temperature. Then, using 1% Triton X 100/PBS, the culture
is passed through a membrane.

To prevent non-specific binding of the antibody,
10 the cells are incubated in 10% goat serum-PBS for 1 hour
and, then, using an anti-tyrosine hydroxylase polyclonal
antibody (Chemicon, diluted 1000-fold in PBS), further
incubated at 4°C for 16 hours. After removal of the
antibody fluid, the cells are washed with PBS and, after
15 addition of peroxidase-labeled dextran polymer-coupled
goat anti-rabbit immunoglobulin (Dako), are incubated at
room temperature for 1 hour.

Detection of the tyrosine-hydroxylase-positive
cells can be made by the color reaction using
20 diaminobenzidine as the substrate. In this manner, the
dopaminergic neuronal survival-supporting activity of the
polypeptide of the invention can be assayed using the
number of tyrosine hydroxylase-positive cells as the
indicator.

25 3) Nerve elongating action

The determination of the nerve elongating action (axonal elongation-promoting action) of the polypeptide of the invention can be carried out using PC12 cells [ATCC Accession Number CRL1721; Science, 229, 393-395 (1985)] as follows. Thus, PC12 cells subcultured in modified Dalbecco's MEM (D-MEM) containing 5% of heat-inactivated (56°C, 30 min) horse serum and 10% fetal calf serum (FCS) are transplanted in a collagen-coated plastic petri dish, 35 mm in diameter, at a concentration of 6×10^4 cells/3 ml. On day 2 after transplantation, the medium was replaced with D-MEM containing a varying concentration of the polypeptide of the invention as well as nerve growth factor (NGF; Wako Pure Chemical Ind.) and FCS and the cultivation is continued in each case. On day 3, morphological changes of the cells are examined with a phase-contrast microscope. By assessing whether the formation of neurites or the promotion of neurite outgrowth is observed in comparison with control, the axonal elongation-promoting potential of the polypeptide of the invention can be evaluated.

4) Neuroglia-activating action

The neuroglia-activating action can be evaluated, for example by determining the effect of the polypeptide of the invention on the activation of neuroglia by FGF in accordance with the method of Kniss et al. or the method

of Bogler et al. [Kniss, D. A., and Burry, R. W., Brain Res., 439, 281-288 (1988); Bogler, O., et al., Proc. Natl. Acad. Sci., USA., 87(16), 6368-6372 (1990)].

5) Mnemonic action

5 The mnemonic action can be evaluated, for example, in accordance with the water-maze protocol of Morris [Morris. R. G. M., J. Neurosci. Meth., 11, 47-60 (1984)].

Another evaluation method comprises administering the LY6H protein or an agonist or antagonist of the LY6H
10 protein as selected by a screening to an animal model of Alzheimer's disease such as a mutant β -amyloid precursor protein gene or mutant presenilin 1 gene transgenic mouse [e.g. Nature, 373, 523-527 (1995); Nature Med., 5, 560-564 (1999)] and evaluating the degree of progression of
15 the disease or the degree of nerve degeneration in comparison with a non-treated control group.

Moreover, in order to have the gene expressed in the human temporal lobe (gene therapy), an adenovirus vector [Straus, E. S., Plenum Press New York, 451-496
20 (1984); Setoguchi, Y., et al., Blood, 84, 2953-2964 (1994)], for instance, is used. Thus, a possible procedure comprises cloning the gene of the invention in an adenovirus vector, culturing it in the stem cell, administering it directly into the temporal lobe or
25 intravenously through a peripheral blood vessel and

checking to see whether Alzheimer type dementia or Alzheimer's disease has been improved or its progression inhibited.

The polypeptide as the active ingredient of the pharmaceutical composition of the invention includes its pharmaceutically acceptable salt. Such salt includes nontoxic salts with alkali metals, alkaline earth metals or ammonium, such as salts with sodium, potassium, lithium, calcium, magnesium, barium and ammonium. These salts can be prepared by the conventional methods in the art. Furthermore, said salt includes nontoxic acid addition salts which can be prepared by reacting the active ingredient polypeptide of the invention with suitable organic or inorganic acids. The representative nontoxic acid addition salt includes the hydrochloride, hydrobromide, sulfate, bisulfate, acetate, oxalate, valerate, oleate, laurate, borate, benzoate, lactate, phosphate, p-toluenesulfonate (tosylate), citrate, maleate, fumarate, succinate, tartrate, sulfonate, glycolate, maleate, ascorbate, benzenesulfonate, naphthalenesulfonate, and the like.

The pharmaceutical composition of the invention includes a composition comprising a pharmacologically effective amount of the polypeptide of the invention and a suitable nontoxic pharmaceutical carrier or diluent.

The pharmaceutical carrier which can be used for said pharmaceutical composition (pharmaceutical preparation) includes diluents or excipients which are conventionally utilized according to dosage forms, such as fillers, volume builders, binders, humectants, disintegrators, surfactants, and lubricants. These can be judiciously selected and used according to the unit dosage form of the composition.

The particularly preferred pharmaceutical composition of the invention can be prepared using various additives which can be formulated in ordinary protein preparations, such as the stabilizer, biocide, buffer, isotonizing agent, chelating agent, pH control agent and surfactant.

The stabilizer includes human serum albumin, an L-amino acid, a sugar, and a cellulose derivative, for instance, can be mentioned. These may be used singly or in combination with a surfactant or the like where necessary. The use in combination with a surfactant may lead to a more effective stabilization of the active ingredient in particular.

The L-amino acid is not particularly restricted but may for example be any of glycine, cysteine and glutamic acid.

The sugar is not particularly restricted but

includes monosaccharides such as glucose, mannose, galactose, fructose, etc.; sugar alcohols such as mannitol, inositol, xylitol, etc., disaccharides such as sucrose, maltose, lactose, etc.; polysaccharides such as dextran, hydroxypropylstarch, chondroitin sulfate, hyaluronic acid, etc.; and their derivatives.

The surfactant is not particularly restricted, either, but both ionic and nonionic surfactants can be employed. Examples of the surfactant are polyoxyethylene glycol sorbitan alkyl esters, polyoxyethylene alkyl ethers, sorbitan monoacyl esters and fatty acid glycerides.

The cellulose derivative that can be used is not particularly restricted, either, but includes methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose, and carboxymethylcellulose sodium.

The level of addition of any of said sugar and other additives can be judiciously selected with reference to the amount in common use. Generally, the sugar is used in a proportion of not less than about 0.0001 mg, preferably within the range of about 0.01 to about 10 mg, per μ g of the active ingredient. The surfactant is used generally in a proportion of not less than about 0.00001 mg, preferably within the range of

about 0.0001 to about 0.01 mg, per μ g of the active ingredient. Human serum albumin, an example of the stabilizer, can be used in a proportion of not less than about 0.0001 mg, preferably within the range of about

5 0.001 to about 0.1 mg, per μ g of the active ingredient. The amount of the amino acid, another example of the stabilizer, can be selected from the range of about 0.001 to about 10 mg per μ g of the active ingredient. The level of addition of the cellulose derivative is not less than

10 about 0.00001 mg and is preferably selected from the range of about 0.001 to about 0.1 mg.

The amount of the active ingredient in the pharmaceutical composition of the invention can be liberally selected from a broad range but is generally

15 selected from the range of about 0.00001 to about 70 weight %, preferably about 0.0001 to about 5 weight %.

The pharmaceutical composition of the invention may be further supplemented with a buffer, an isotonizing agent, and a chelating agent. The buffer includes boric

20 acid, phosphoric acid, acetic acid, citric acid, ϵ -aminocaproic acid, glutamic acid, and the corresponding salts (the alkali metal or alkaline earth metal salts thereof, such as sodium salts, potassium salts, calcium salts and magnesium salts). The isotonizing agent

25 includes sodium chloride, potassium chloride, sugars and

glycerol. The chelating agent includes sodium edetate, and citric acid. The level of addition of any of these additives may be within the conventional range.

The pharmaceutical preparation of the invention can
5 be provided in the form of a solution, and in a lyophilized form which can be stored. Such lyophilized preparations can be extemporaneously dissolved in, for example, a buffer inclusive of water, saline or the like at a suitable concentration.

10 As regards the unit dosage form of the pharmaceutical composition of the invention, various forms can be selected according to the therapeutic objective. The representative form includes solid dosage forms such as tablets, pills, powders, neat powders, granules, capsules,
15 etc. and liquid dosage forms such as solutions, suspensions, emulsions, syrups, elixirs and so on. These dosage forms are generally classified, by route of administration, into oral preparations, parenteral preparations, nasal preparations, vaginal suppositories, rectal
20 suppositories, sublingual tablets, ointments, and others. Each of such dosage forms can be formulated and molded or otherwise prepared by the established pharmaceutical procedure.

For example, tablets can be manufactured using, as
25 said pharmaceutical carrier, any of various excipients

such as lactose, sucrose, sodium chloride, glucose, urea, starch, calcium carbonate, kaolin, crystalline cellulose, silicic acid, potassium phosphate, etc.; binders such as water, ethanol, propanol, simple syrup, glucose solution, starch solution, gelatin solution, carboxymethylcellulose, hydroxypropylcellulose, methylcellulose, polyvinylpyrrolidone, etc.; disintegrators such as carboxymethylcellulose sodium, carboxymethylcellulose calcium, low-substitution-degree hydroxypropylcellulose, dried starch, sodium alginate, agar powder, laminaran powder, sodium hydrogen carbonate, calcium carbonate, etc.; surfactants such as polyoxyethylene sorbitan fatty acid esters, sodium lauryl sulfate, monoglycerol stearate, etc.; disintegration inhibitors such as sucrose, stearin, cacao butter, hydrogenated oil, etc.; absorption promoters such as quaternary ammonium bases, sodium lauryl sulfate, etc.; humectants such as glycerol, starch, etc.; adsorbents such as starch, lactose, kaolin, bentonite, colloidal silica, etc.; and lubricants such as purified talc, salts of stearic acid, boric acid powder, polyethylene glycol and so on.

Where necessary, such tablets can be coated with conventional coatings to provide sugar-coated tablets, gelatin-coated tablets, enteric-coated tablets and film-coated tablets. Double-layer or multi-layer tablets may

also be employed.

The pharmaceutical carrier which can be used for the production of pills includes various excipients such as glucose, lactose, starch, cacao butter, hydrogenated
5 vegetable oil, kaoline, talc, etc.; binders such as gum arabic powder, tragacanth powder, gelatin, ethanol, etc.; and disintegrators such as laminaran and agar.

The capsules can be generally manufactured in the conventional manner by blending the active ingredient of
10 the invention with the pharmaceutical carrier or carriers and filling the resulting composition into hard gelatin capsule shells, soft capsule shells or the like.

The liquid preparation for oral administration includes pharmaceutically acceptable solutions, emulsions,
15 suspensions, syrups, elixirs, etc. as formulated with routine inert diluents, such as water, and these dosage forms may contain a wetting agent, an emulsifier, a suspending agent and/or other auxiliary additives. These can be manufactured by the established pharmaceutical
20 procedures.

The liquid preparation for parenteral administration, inclusive of sterile aqueous and non-aqueous solutions, emulsions and suspension, can be prepared using such diluents as water, ethyl alcohol, propylene
25 glycol, polyethylene glycol, ethoxylated isostearyl

alcohol, polyoxylated isostearyl alcohol, polyoxyethylene sorbitan fatty acid esters, and vegetable oils such as olive oil. In addition, injectable organic esters, such as ethyl oleate, may be formulated. Furthermore, any of
5 the conventional solubilizers, buffers, wetting agents, emulsifiers, suspending agents, preservatives, dispersants, etc. can also be added.

The above various pharmaceutical dosage forms are sterilized in the routine manner. This sterilization can
10 be achieved by filtration through a bacterial filter, formulation of a biocide, irradiation, or a heat treatment. Furthermore, these may be provided in the form of sterile solid compositions which can be extemporaneously dissolved in sterile water or a suitable
15 sterilizable medium.

For the manufacture of dosage forms for rectal or vaginal administration, such pharmaceutical carriers as polyethylene glycol, cacao butter, higher alcohols, higher alcohol esters, gelatin, semisynthetic glycerides,
20 etc. can be employed.

Ointments such as pastes, creams and gels can be prepared using a diluent such as white petrolatum, paraffin, glycerol, cellulose derivatives, propylene glycol, polyethylene glycol, silicone, bentonite, and
25 vegetable oils such as olive oil.

Compositions for transnasal or sublingual administration can be prepared in the routine manner using a well-known standard excipient.

Where necessary, those pharmaceutical preparations
5 of the invention may be supplemented with coloring agents, preservatives, perfumes, flavoring agents, sweeteners, and other drugs.

The method for administration of such pharmaceutical preparations is not particularly restricted but can
10 be selected according to the specific dosage form, patient's age, sex and other factors, severity of illness, and other variables. For example, tablets, pills, solutions, suspensions, emulsions, granules and capsules are administered orally, while parenteral products are
15 administered intravenously, either alone or in admixture with the conventional glucose, amino acid or other infusion, or, where necessary, administered alone intramuscularly, intradermally, subcutaneously or intraperitoneally. The rectal suppositories are administered into
20 the rectum; the vaginal suppositories are administered into the vagina; the nasal preparations are administered into the nostrils, the sublingual preparations are administered buccally, and the ointments are administered topically for transdermal drug delivery.

25 The dosage for any of the above pharmaceutical

preparations is not particularly restricted but can be judiciously selected from a broad range according to the expected therapeutic effect, administration method, duration of treatment, patient background such as age and sex, and other factors. Generally, the recommended usual dosage of the active ingredient is about 0.01 μ g - 10 mg/day, preferably about 0.1 μ g - 1 mg/day, per kg of the patient's body weight. The above dose may be administered once a day or in 2 or more divided doses.

Furthermore, as pointed out in the working example to be presented later herein, the expression of the gene of the invention has been abolished or decreased in the temporal lobe of patients with Alzheimer's disease. Therefore, by constructing an arbitrary expression vector harboring the whole or part of the gene of the invention and introducing the expression vector into the temporal lobe tissue for forced expression of the gene in the tissue, neurodegenerative changes inclusive of an excessive atrophy of neurons in the temporal lobe may be inhibited and, hence, the progression of Alzheimer's disease may be arrested. Therefore, the present invention further provides a pharmaceutical composition for gene therapy (gene therapeutic agent) which is possessed of such a neurodegeneration-inhibitory action.

The present invention further provides the above

expression vector or vector for gene therapy, cells
transfected with the gene of the invention through the
introduction of said vector, and a pharmaceutical
composition for gene therapy which comprises any of the
5 above as the active component.

The gene therapy using said gene therapeutic agent
is performed by administering at least one member
selected from the group consisting of the vector for
introduction and expression of the gene of the invention
10 and cells transfected with the gene of the invention
through the introduction of said vector into the brain
neurons or temporal lobe tissue of a patient with
neurodegenerative disease. By such a procedure,
neurodegenerative changes in such tissue can be inhibited
15 and symptoms of Alzheimer's disease, Alzheimer type
dementia, Parkinson's disease, brain ischemia, etc. can
be alleviated.

The gene therapy is now described in further detail.
In the following execution of a gene therapy, the routine
20 chemical, molecular biological, microbiological,
recombinant DNA, genetic, and immunological techniques
can be employed unless otherwise specified. These
techniques are described in, inter alia, Maniatis, T.,
et al., Molecular Cloning: A laboratory manual (Cold
25 Spring Harbor Laboratory), Cold Spring Harbor, New York

(1982)), Sambrook, J., et al., Molecular Cloning: A laboratory manual, 2nd Ed. (Cold Spring harbor Laboratory), Cold Spring harbor, New York (1981)), Ausbel, F. M., et al., Current Protocols in Molecular Biology, 5 John Wiley and Sons, New York (1992)), Glover, D., DNA Cloning, I and II (Oxford Press (1985)), Anand, Techniques for the Analysis of Complx Genomes (Academic Press (1992)), Guthrie, G., et al., Guide to Yeast Genetics and Molecular Biology (Academic Press (1991)), 10 and Fink, et al., Hum. Gene Ther., 3, 11-19 (1992).

The gene therapy can be carried out using a gene therapy vector harboring the whole or part of the gene of the invention or cells transfected with the gene of the invention through the introduction of said vector. This 15 gene therapy may for example be a method of supplying the LY6H gene or its function to cells in which said gene has not been expressed. By such gene therapy, neurodegeneration around the receptor cell/target cell is inhibited.

20 The gene of the invention or a fragment of the gene can be introduced into cells by means of a vector adapted to maintain the gene extrachromosomally. In such cases, the particular gene can be caused to be expressed by the cells from an extrachromosomal position. Moreover, when 25 the LY6H gene is to be expressed by introducing a

fragment of the gene into the temporal lobe site of the brain nervous system where no expression of the gene is found, the particular fragment of the gene may be a fragment encoding a part of LY6H protein which is
5 necessary for the survival or non-tumorigenic growth of cells.

The gene transfer vector may be any of various known vectors in which the gene of the invention has been subcloned as will be described later herein.

10 The introduction of the gene transfer vector into the target cell can be easily effected by the established technology of introducing DNA into various cells which is already known to those skilled in the art, such as electroporation, calcium phosphate transfection
15 (coprecipitation), virus transduction and other techniques. The cells transfected with the gene of the invention can be utilized as a drug for neurodegenerative disorders of the brain, inclusive of an inhibitor of premature atrophy of the brain nervous system, or as
20 models for therapeutic research.

As mentioned above, the gene or gene fragment of the invention as introduced by the gene therapy according to the invention increases the expression of the corresponding gene product in the brain nerve or
25 surrounding tissue to thereby inhibit atrophy of the

brain nerve in the tissue expressing the gene. Such gene therapy can be applied with advantage to the brain neuronal tissue where the expression of LY6H gene or the LY6H protein has been abolished as well as to the brain
5 neuronal tissue where the level of expression of said gene has been depressed.

The gene therapy according to the present invention is performed as follows. First, a screening is carried out for candidate patients for the gene therapy by
10 recording a computer tomogram (CT) with the scanning position fixed to the temporal lobe of the patient with Alzheimer type dementia or Alzheimer's disease to check for atrophy of the temporal lobe or progression of the atrophy.

15 Then, to achieve expression of the gene of the invention, the intracellular LY6HmRNA is created in the target cell and its translation is promoted to accelerate expression of the LY6H gene. For this purpose, preferably a sense oligonucleotide corresponding to the mRNA of the
20 gene is produced and supplied to the target cell. By providing the cell with the activity to promote expression of the LY6H gene by the above gene therapy, the neurodegenerative change in the brain receptor cell/target cell can be inhibited.

25 According to the above gene therapy using said

product in the target cell can be used as such vectors. Any of the expression vectors mentioned above can be usually employed but the preferred are vectors constructed by using any of the vectors disclosed in USP 5 5252479 and WO 93/07282 (specifically pWP-7A, pWP-19, pWU-1, pWP-8A, pWP-21 and/or pRSVL) or the pRC/CMV (Invitrogen) as the source vector. The still more preferred are the various virus vectors described later herein.

10 As the promoter for use in the vector for gene therapy, those promoters, which are intrinsic to the affected tissues to be treated in various diseases, are preferably employed. Examples of the promoters are albumin, α -fetoprotein, α 1-antitrypsin, transferrin, and 15 transthyretin for the liver, and carbonic anhydrase I and carcinoembryonic antigen for the colon. When the affected tissues are the uterus and placenta, estrogen, aromatase, cytochrome P450, cholesterol side-chain-cleaving enzyme P450, and 17 α -hydroxylase P450 can be exemplified.

20 For the prostate, prostate-specific antigens, gp91-fox gene, and prostate-specific kallikrein can be exemplified. For the breast, erb-B2, erb-B3, β -casein, β -lactoglobulin, and whey protein can be exemplified. For the lung, surfactant protein C, and uroglobulin can be 25 exemplified. For the skin, K-14-keratin, human keratin 1

or 6, and leucine can be exemplified. For the brain,
glial fibrillary acidic protein, mature astrocyte-
specific protein, myelin, tyrosine hydroxylase pancreatic
villin, glucagon, and Langerhans islet amyloid
5 polypeptide can be exemplified. For the thyroid,
thyroglobulin, and calcitonin can be exemplified. For the
bone, $\alpha 1$ collagen, osteocalcin, and bone
sialoglycoprotein can be exemplified. For the kidney,
renin, liver/bone/kidney alkaline phosphatase, and
10 erythropoietin can be exemplified, and for the pancreas,
amylase, and PAP1 can be exemplified.

Furthermore, in the production of a vector for
introduction of a sense oligonucleotide, the sense
oligonucleotide to be introduced (one having a full-
15 length or partial sequence corresponding to the sequence
of the gene of the invention) can be easily prepared and
acquired by the standard genetic engineering techniques
based on the nucleotide sequence information on the gene
of the invention as described hereinbefore.

20 The transfer of such a vector for introduction of a
sense oligonucleotide into cells can be carried out by
various techniques already known in the art, such as
electroporation, calcium phosphate transfection
(coprecipitation), virus transduction and the like. The
25 cells transfected with said sense oligonucleotide, as

sense oligonucleotide, the objective inhibition of neurodegenerative change of the brain and consequent alleviation or arrest of progression of neurodegenerative symptoms may be successfully attained by subcloning the
5 LY6H gene into a retrovirus, adenovirus or AAV-derived vector and infecting the target brain nerve cells with the vector to thereby cause expression of the sense oligonucleotide.

When a sense oligonucleotide of the gene of the
10 invention is introduced into the cerebral neuron or tissue to increase the expression of LY6H protein, the sense oligonucleotide need not be the full-length nucleotide of the LY6H gene but may be the modification product insofar as it retains a function substantially
15 identical to the function of the parent gene and promotes expression of the LY6H gene or a fragment gene comprising a partial sequence retaining said function.

Vectors which can be used for introducing an objective gene for both DNA recombination and
20 extrachromosomal gene maintenance are already known in the art and any of such known vectors can be used in the practice of the invention. For example, a virus or plasmid vector which includes a copy of LY6H gene sense oligonucleotide ligated to the expression control element
25 and is capable of expressing the sense-oligonucleotide

such and in an isolated form, have a brain neurodegeneration-inhibitory action so that they can be used as a drug, or a therapeutic research model, for the inhibition or arrest of progression of neurodegenerative
5 lesions as well.

In gene therapy, the above vector for introduction of a sense oligonucleotide can be injected either topically into the temporal lobe or surrounding region of the patient or systemically. Furthermore, it may be
10 cultured together with stem cells and, then, administered by local or systemic injection. By such administration, the vector can be introduced into the nerve cells of the patient's brain. In the event the transduced gene is not permanently taken up in the chromosome of each target
15 cell, the administration may be repeated periodically.

The method for gene therapy according to the invention includes both the in vivo technique which comprises administering a construct for introduction of said sense oligonucleotide (a sense oligonucleotide
20 transfer vector) directly into the body and the ex vivo technique which comprises transferring the gene into cultured stem cells and, after culturing, transplanting or otherwise introducing the cells into the patient's body. A gene therapy comprising introducing said sense
25 oligonucleotide directly into the cell is also feasible.

The target cells into which the sense oligonucleotide of the gene of the invention is to be introduced can be judiciously selected according to the object of gene therapy (treatment). For example, the
5 target cells include brain neurons and brain nerve tissues as well as lymphocytes, fibroblasts, hepatocytes and hemopoietic cells.

The method of introducing the sense oligonucleotide in the above gene therapy includes a viral introduction
10 technique and a non-viral introduction technique.

As to the viral introduction technique, in consideration of the fact that the sense oligonucleotide to be transferred is a foreign substance which is expressed especially in the normal brain cells, the
15 method using a retrovirus vector, for instance, can be exemplified. Other virus vectors which can be used include the adenovirus vector, HIV (human immunodeficiency virus) vector, adeno-associated virus (AAV) vector, herpes virus vector, herpes simplex virus
20 (HSV) vector, and Epstein-Barr virus (EBV) vector.

The method of constructing a virus vector for transfer of a sense oligonucleotide and the method for transfer of the sense oligonucleotide to the target cell or target tissue are now specifically described.

25 The retrovirus vector system consists of a virus

vector and a helper cell (packaging cell). The helper cell means a cell which has expressed genes encoding the structural protein gag (structural protein within the virus particle), pol (reverse transcriptase), env (coat protein), etc. of a retrovirus but which has not formed virus particles. On the other hand, the virus vector has the packaging signal and LTR (long terminal repeats) but lacks structural genes, such as gag, pol, env, etc., which are necessary for virus replication. The packaging signal is a sequence which functions as a tag in the assembly of a virus particle. Selective genes (neo, hyg) and the object sense oligonucleotide ligated in the cloning site are inserted in lieu of the virus genes. In order that a high titer of virus particles may be obtained, it is important to use an insert as short as possible, provide a broad packaging signal including a part of the gag gene, and use care not to leave ATG of the gag gene.

As the vector DNA harboring the object sense oligonucleotide is transferred to the helper cell, the vector genomic RNA is packaged by the virus structural protein formed by the helper cell, whereby virus particles are formed and secreted. The virus particle as a recombinant virus infects the target cell and, as a result, the DNA sequence reverse-transcribed from the

virus genomic RNA is integrated into the cell nucleus so that the sense gene inserted in the vector is expressed.

It may be employed a technique using a fibronectin fragment containing the cell adhesion domain, heparin-
5 binding site and conjugating segment [Hanenberg, H., et al., Exp. Hemat., 23, 747 (1995)], for enhancing the efficiency of transfer of the object gene.

An example of the retrovirus vector for use in the above retrovirus vector system is the retrovirus derived
10 from mouse leukemia virus [McLachlin, J. R., et al., Proc. Natl. Acad. Res. Molec. Biol., 38, 91-135 (1990)].

The method using an adenovirus vector is now described in detail. The adenovirus vector can be constructed in accordance with the methods described in
15 Berkner, K. L., Curr. Topics Microbiol. Immunol., 158, 39-66 (1992), Setoguchi, Y., et al., Blood, 84, 2946-2953 (1994), Kanegae, H. et al. [Jikken Igaku (Experimental Medicine), 12, 28-34 (1994)] and Ketner, G. et al., Proc. Natl. Acad. Sci., USA., 91, 6186-6190 (1994).

20 For example, to construct a non-proliferative adenovirus vector, the early region E1 and/or E3 of the adenovirus is excised in the first place. Then, a plasmid vector containing the desired foreign gene expression unit (which consists of the sense oligonucleotide to be
25 transferred, the promoter for transcription of said sense

oligonucleotide, Poly A for insuring the stability of the transcribed gene) and a part of the adenovirus genomic DNA and a plasmid containing the adenovirus genome are used to cotransfect the 293 cell, for instance. As a
5 homologous recombination is thus caused to take place between them for substitution of the gene expression unit for E1, a nonproliferative adenovirus vector is obtained as a vector harboring the object sense oligonucleotide. A
3'-end adenovirus vector with a terminal protein added
10 can also be constructed by ligating the adenovirus genomic DNA in a cosmid vector. Furthermore, the YAC vector may also be utilized for the construction of an adenovirus vector.

Production of an adeno-associated virus (AAV)
15 vector is now described briefly. AAV was discovered as a small virus contaminating adenovirus culture systems. As to this virus, the existence of the genus Parvovirus capable of autonomous proliferation within the host cell without requiring a helper virus for virus replication
20 and the genus Dependovirus which requires a helper virus has been identified. This AAV has a broad host range and is one of the common viruses infecting various kinds of cells. The virus genome is a linear single-stranded DNA consisting of 4680 nucleotides, with the 145 nucleotides
25 at both ends having a characteristic sequence known as

ITR (inverted terminal repeat). This ITR region functions as the replication origin and plays the role of a primer. This ITR is also essential to packaging for virus particles and integration of AAV into the chromosome DNA of the host cell. In regard of the virus protein, the left-half of the genome codes for the nonstructural protein, that is the regulatory protein Rep which controls replication and transcription.

Construction of the recombinant AAV can be carried out by utilizing the property of AAV to become integrated into the chromosome DNA, whereby the desired gene transfer vector can be prepared. This method may be described in detail as follows. First, a plasmid (AAV vector plasmid) retaining the ITRs at 5'- and 3'-ends of a wild-type AAV and harboring the sense oligonucleotide to be transferred as interposed therebetween is constructed. The virus protein necessary for virus replication and formation of virus particles is supplied from a separate helper plasmid. It is necessary to insure that no common nucleotide sequence will exist between the two plasmids so that a wild-type virus will not appear on DNA recombination. Thereafter, the two plasmids are transferred into the 293 cell by transfection, for example, and, further, the cells are infected with an adenovirus as the helper virus (when the 293 cell is used,

this adenovirus may be a non-proliferative one), whereby the desired non-proliferative recombinant AAV is produced. Since this recombinant AAV is present in the nucleus, the cells are subjected to freeze-thawing and recovered and
5 the contaminant adenovirus is inactivated by heating at 56°C. Then, where necessary, the recombinant AAV is separated and concentrated by ultracentrifugation using cesium chloride. In this manner, the desired recombinant AAV for gene transfer can be obtained.

10 Production of an EBV vector can be carried out by the method of Shimidzu et al. [Shimidzu, N., SAIBO KOUGAKU (Cell Technology, 14(3), 280-287 (1995))].

 Production of the EBV vector for transfer of the sense oligonucleotide according to the invention is now
15 described briefly. EB virus (Epstein-Barr virus) is a virus of the family Herpesviridae, which was first isolated by Epstein and coworkers from cultured cells derived from Burkitt lymphoma [Kieff, E. and Liebowitz, D.: Virology, 2nd ed. Raven Press, New York, 1990, pp.
20 1889-1920]. This EBV has cell-transforming activity and in order to use it as a vector for gene transfer, it is necessary to prepare a virus defected of this transforming activity. This can be done as follows.

 Thus, first of all, the EBV genome in the vicinity
25 of the target DNA in which the desired foreign gene is to

be inserted is cloned. Then, a DNA fragment of the foreign gene and a drug-resistant gene are inserted to construct a vector for preparation of a recombinant virus. Then, the vector for preparation of a recombinant virus
5 as excized with a suitable restriction enzyme is transfected to EBV-positive Akata cells. The recombinant virus formed by homologous recombination is recovered, together with the wild type Akata EBV, through stimulation of virus production by anti-surface
10 immunoglobulin treatment. The recombinant virus is infected to EBV-negative Akata cells and, in the presence of a drug, resistant clones are selected, whereby Akata cells infected exclusively with the recombinant virus free of wild type EBV can be obtained. Further, by
15 inducing viral activity in the recombinant virus-infected Akata cells, the objective recombinant virus vector can be produced in quantities.

The method of introducing the object gene into the target cell or target tissue in the gene therapy of the
20 invention includes the following representative two methods.

The first method comprises harvesting the target cells from a patient to be treated, growing the cells ex vivo, for example under addition of interleukin-2 (IL-2)
25 or the like, to transfer the objective sense

oligonucleotide harbored in the retrovirus vector, and retransplanting the resulting cells (ex vivo method). This method is suitable for the therapy of genetic diseases caused by defective genes and cancer, for instance.

The second method is a method for direct gene transfer which comprises injecting the object sense oligonucleotide directly into the patient's body or the target site such as the cerebral tissue (direct method).

More particularly, the first method can be carried out in the following manner, for instance. Thus, the mononuclear cells, such as stem cells, harvested from the patient are fractionally separated from monocytes using a blood sorter and cultured in the presence of IL-2 in a suitable medium such as AIM-V medium for about 72 hours, followed by addition of the vector harboring the sense oligonucleotide to be introduced. For enhancing the efficiency of transfer of the sense oligonucleotide, the cells may be grown in the presence of protamine at 32°C for 1 hour, centrifuged at 2500 ppm, and then cultured under 10% carbon dioxide gas at 37°C for 24 hours. After this procedure is repeated a few times, the cells are further cultured in the presence of IL-2 in, for example, AIM-V medium for 48 hours and then washed with saline.

The viable cells are counted and the efficiency of

introduction of the sense oligonucleotide is evaluated by said in situ PCR or, when the object is enzymatic activity, assaying the degree of the enzymatic activity.

The safety checks such as culture of bacteria and
5 fungi in cultured cells, check for the presence or absence of mycoplasma infection, search for endotoxin, etc. are carried out to confirm safety. Thereafter, the cultured cells transformed with the predicted effective
10 dose of the sense oligonucleotide are returned to the patient by intravenous drip injection. The above procedure is repeated at intervals of several weeks or a few months to consummate the gene therapy.

The dosage of the virus vector is judiciously selected according to the target cell. The usually
15 preferred dose may for example be 1×10^3 cfu - 1×10^8 cfu in terms of virus titer per 1×10^8 target cells.

It can be adopted an alternative version of the above first method that comprises co-cultivating the virus-producer cells having the retrovirus vector
20 harboring the object sense oligonucleotide and the patient's cells to thereby introduce the sense oligonucleotide into the target cells.

In carrying out the second method (direct method) for gene therapy, it is particularly preferable to
25 perform a preliminary experiment ex vivo to check whether

the objective sense oligonucleotide can be actually introduced by the gene transfer method by carrying out PCR of the vector gene cDNA or in situ PCR or check whether the desired therapeutic effect, for example

5 elevation of a specific activity or the growth or inhibition of growth of the target cell can be actually achieved by introduction of the objective sense oligonucleotide. Moreover, when a virus vector is used, it is, of course, of great importance to confirm the

10 safety of introduction of the sense oligonucleotide in gene therapy by performing a PCR search for proliferative retrovirus and the like, determining the reverse transcriptase activity, or monitoring the coat protein (env) gene by the RCR technique.

15 The gene therapy of the invention in Alzheimer's disease, Alzheimer type dementia or Parkinson's disease may for example be a therapy of neurodegenerative disease which comprises harvesting stem cells or brain nerve cells from the patient, establishing a cultured cell line

20 by enzymatic treatment or the like, introducing the object sense oligonucleotide into the target brain nerve cells utilizing AAV or the like, carrying out a screening with G418 cells, measuring the amount of expression of IL-12 or the like in vivo, giving a radiation treatment,

25 and inoculating the cells into the patient's brain tissue

or the temporal lobe site.

The present invention further provides a pharmaceutical composition or preparation (a gene therapeutic agent) comprising a sense oligonucleotide transfer vector of the invention or a cell line transformed with the sense oligonucleotide as an active ingredient in a pharmacologically effective amount in combination with a suitable nontoxic pharmaceutical carrier or diluent.

10 The pharmaceutical carrier that can be utilized in the pharmaceutical composition (pharmaceutical preparation) of the invention includes those diluents or excipients, e.g. fillers, volume builders, binders, humectants, disintegrators, surfactants, lubricants, etc.,
15 which are usually employed depending on the mode of use of such a composition, and these can be selectively used according to the contemplated unit dosage form of the preparation.

 The unit dosage form of the pharmaceutical
20 preparation of the invention may be the same as mentioned for the polypeptide preparation of the invention, and a suitable one can be judiciously selected according to the therapeutic objective.

 The therapeutic and prophylactic method for
25 neurodegenerative disease according to the invention is

now described in detail.

The present invention provides a method for therapy of neurodegenerative diseases, such as Alzheimer's disease, Alzheimer type dementia, brain ischemia, Parkinson's disease, and like diseases in which either an excess or a shortage of the LY6H polypeptide is involved. When LY6H activity is excessive, several approaches can be taken. The first method comprises administering an inhibitor compound (antagonist) in an effective amount to inhibit the function of LY6H polypeptide by blocking its binding to a ligand, substrate, receptor, enzyme or the like or inhibiting a secondary signal in combination with a pharmaceutically acceptable carrier to thereby improve an abnormal state. An alternative method comprises administering a soluble-form LY6H polypeptide capable of binding to a ligand, substrate, enzyme, receptor or the like in competition with the endogenous LY6H. A typical example of such competitive substance includes a fragment of LY6H polypeptide. In another method, a soluble-form LY6H polypeptide capable of binding to a ligand in competition with endogenous LY6H can be administered. A typical example of such competitive substance includes a fragment of LY6H polypeptide.

In a still another method, expression of the gene coding for endogenous LY6H polypeptide can be inhibited

by applying a gene expression inhibition technique to the LY6H gene product. The known technique of this kind includes the use of an internally generated or separately administered antisense sequence [e.g. Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988), O'Connor, J. Neurochem 56: 560 (1991)]. As an alternative method, an oligonucleotide capable of forming a triple helix with the gene can be supplied [e.g. Lee et al., Nucleic Acids Res., 6: 3073 (1979); Cooney et al., Science, 241: 456 (1988); Dervan et al., Science, 251: 1360 (1991)]. These oligomers can be administered as such or related oligomers may be caused to be expressed in vivo.

For the therapy of abnormal symptoms related to an under-expression of LY6H and its activity, several methods can be utilized. The first method comprises administering a compound capable of activating LY6H (agonist) in a therapeutically effective amount together with a pharmaceutically acceptable carrier to a subject to thereby improve the abnormal symptoms. In another method, the endogenous production of LY6H by related cells in the subject can be actuated by gene therapy. For example, the polynucleotide of the present invention may be manipulated so as to be expressed with a defective retrovirus vector as mentioned hereinbefore. Then, this

retrovirus expression construct is isolated and introduced into packaging cells transduced with a retrovirus plasmid vector harboring the RNA encoding the polypeptide of the invention so that the packaging cells will form infective virus particles containing the object gene. These producer cells are administered to the subject for in vivo manipulation of the cells so that the polypeptide may be expressed in vivo. For an overview of gene therapy, reference may be made to Human Molecular Genetics, T. Strachan and A. P. Read, BIOS Scientific Publishers Ltd. (1996), Chapter 20 - Gene Therapy and Other Molecular Genetic-based Therapeutic Approaches, inclusive of the specific references cited therein. An alternative method comprises administering a therapeutic dose of LY6H polypeptide in combination with a suitable pharmaceutical carrier.

The cells may for example be formulated in phosphate-buffered saline (pH 7.4), Ringer's solution or an intracellular composition injection or in such a dosage form as can be administered in combination with a substance conducive to an enhanced gene transfer efficiency, such as protamine.

The method of administering the above pharmaceutical preparation is not particularly restricted but a suitable regimen can be established according to the

particular dosage form, the patient's age, sex and other factors, the severity of illness, and the like.

The amount of the active ingredient to be incorporated in the pharmaceutical preparation and the dosage are not particularly restricted but each can be
5 liberally selected from a broad range according to the expected therapeutic benefit, method of administration, duration of treatment, patient background inclusive of age and sex, and other variables.

10 Generally, the dosage of the retrovirus vector harboring the sense oligonucleotide as a pharmaceutical preparation may for example be about 1×10^3 pfu through 1×10^{15} pfu in terms of retrovirus titer per kilogram body weight per day.

15 In the case of cells carrying the sense oligonucleotide for introduction, the dosage can be properly selected from the range of about 1×10^4 cells/body through 1×10^{15} cells/body.

The above preparation can be administered once a
20 day or in a few divided doses a day, or even intermittently at intervals of 1 or several weeks. Preferably, a substance conducive to an enhanced gene transfer efficiency, such as protamine, or a preparation containing the same can be administered in combination.

25 When the gene therapy according to the invention

is applied to the therapy of a neurodegenerative disease, it can be performed in a suitable combination with other gene therapies (conjunctive gene therapy) or in combination with a pharmacotherapy utilizing an

5 acetylcholinesterase inhibitor or the like and/or a rehabilitation therapy. The gene therapy of the invention can be performed with reference to the NIH guidelines, inclusive of its safety aspect [Recombinant DNA Advisory Committee, Human Gene Therapy, 4, 365-389 (1993)].

10 Furthermore, in accordance with the invention, for the purpose of detecting the presence of LY6H gene, it is possible to prepare a biological sample such as blood or serum, optionally extract the nucleic acid, and analyzing it for LY6H gene.

15 The method of detecting the gene may comprise preparing a DNA fragment of the gene of the invention and design it so that it may be used in the screening for LY6H gene and/or its amplification. More specifically, it is possible to construct a DNA fragment having the
20 properties of a probe for plaque hybridization, colony hybridization, Southern blotting, Northern blotting, etc. or a probe for the preparation of a full-length or partial DNA of the gene of the invention as amplified by a polymerase chain reaction (PCR) which amplifies a
25 nucleotide sequence with a polymerase. For this purpose,

a primer having the same sequence as LY6H gene is first prepared. Then, this primer is reacted, as a probe for screening, with a biological sample (nucleic acid sample) to check for the presence of the particular LY6H gene
5 sequence. The nucleic acid sample may be prepared by any of various techniques facilitating detection of the target sequence, such as denaturation, restriction enzyme digestion, electrophoresis or dot blotting.

As the method for said screening, the use of a PCR
10 technique is particularly preferred from sensitivity points of view, and this technique is not particularly restricted inasmuch as a fragment of the gene of the invention is used as a primer. Thus, It can be utilized that any of the hitherto-known techniques [Science, 230,
15 1350-1354 (1985)] and the modified versions of PCR which have been developed of late or will be developed in the future [Sakaki, Yoshiyuki et al. (ed.), Jikken Igaku (Experimental Medicine), Supplement 8(9) (1990), Yōdoshā; Protein, Nucleic Acid, Enzyme: Special Supplement,
20 Kyoritsu Shuppan, 35(7) (1990)].

The DNA fragment for use as the primer is a chemically synthesized oligo-DNA, and such oligo-DNA can be synthesized using an automated DNA synthesizer or the like, for example Pharmacia LKB Gene Assembler Plus
25 (Pharmacia). The preferred length of the primer (sense

primer or antisense primer) to be synthesized may for
example be about 10-30 nucleotides. The probe for us in
said screening is usually a labeled probe but may be an
unlabeled one, or the detection may be made according to
5 specific binding to a directly or indirectly labeled
ligand. The suitable label and the method of labeling the
probe or ligand belong to the prior art. Thus, the prior
art label includes radioisotopes, biotin, fluorescent
groups, chemiluminescent groups, enzymes, antibodies,
10 etc., which can be taken up through known procedures such
as nick translation, random priming and kinase treatment.

The PCR technique to be used for detection may for
example be RT-PCR but various modifications of the
technique which are in routine use in the art can be
15 utilized.

Furthermore, the above assay method can be
expediently carried out by utilizing an reagent kit for
detecting an LY6H gene in samples.

Therefore, the present invention provides an LY6H
20 gene detection reagent kit comprising a DNA fragment of
the gene of the invention.

This reagent kit comprises at least a DNA fragment
which hybridizes with a part or the whole of the
nucleotide sequence shown in SEQ ID NO:2 or its
25 complementary nucleotide sequence as an essential

component and may optionally contain other components such as a labeling agent and PCR reagents (for example, Taq DNA polymerase, deoxynucleotide triphosphates, primers, etc.).

5 The labeling agent may be a radioisotope or a chemical modifier such as a fluorescent substance but the DNA fragment as such may have been conjugated with such a labeling agent. This reagent kit may further contain a suitable reaction solvent or diluent, standard antibody,
10 buffer, wash solution, reaction stopper solution, etc. which make an assay easier to perform.

 The present invention in a further aspect provides a method for diagnosis of neurodegenerative diseases which comprises using the above assay method and a
15 diagnostic agent or diagnostic reagent kit for use in practicing said method.

 By the direct or indirect sequencing of the LY6H genes obtained from test samples by utilizing the above method, it is possible to find new LY6H gene-related
20 genes having high homology to the wild-type LY6H gene.

 Therefore, the present invention further provides a method of screening for human LY6H gene-related genes in samples which comprises performing said assay and sequencing of the LY6H genes contained in test samples.

25 The wild-type LY6H and/or mutant LY6H can be

determined by utilizing the protein encoded by the human LY6H gene of the invention (a polypeptide having the amino acid sequence shown in SEQ ID NO:1), a polypeptide having an amino acid sequence derived from the sequence shown in SEQ ID NO:1 by the deletion, substitution or addition of 1 or a plurality of amino acids, a fragment of either of them, or an antibody to any of such proteins.

Therefore, the invention provides a method of determining an anti-wild-type LY6H and/or mutant LY6H antibody or a method of determining the antigen. By this method, the degree of impairment of the brain nerve can be detected from a change in wild-type LY6H (polypeptide). Such changes can be detected by the sequencing of LY6H by the well-established technology described hereinabove, more preferably by detecting differences in the LY6H polypeptide or the presence or absence of LY6H polypeptide by the use of said antibody (polyclonal or monoclonal antibody).

The following is a specific example of determination of said wild-type and/or mutant LY6H. The anti-LY6H antibody can be used to immunoprecipitate LY6H polypeptide from a solution containing a biological sample obtained from a human body, such as blood or serum or can be reacted with the LY6H polypeptide on polyacrylamide gel of Western blot or immunoblot. The

LY6H polypeptide in a paraffin section or frozen tissue specimen can be detected by an immunohistochemical technique using the anti-LY6H antibody. The antibody production and purification technology are well known in the art and suitable techniques can be selectively employed.

The preferred technology relevant to the detection of a wild-type LY6H or a mutant thereof includes enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), immunoradiometric assay (IRMA) and immunoenzymometric assay (IEMA) with a sandwich technique using a monoclonal antibody and/or a polyclonal antibody.

The invention further provides an LY6H ligand or an LY6H receptor existing in a cell membrane fraction or on a cell surface and having binding affinity for LY6H polypeptide. The LY6H receptor can be obtained by conjugating a labeled LY6H polypeptide in a biological sample containing a cell membrane fraction, extracting, isolating and purifying the conjugation product and identifying the amino acid sequence of the isolated product. The procedure for preparation and the method of sequencing this LY6H receptor polypeptide is obvious to one skilled in the art.

Furthermore, by applying the LY6H receptor or a fragment thereof to a screening for various drugs, the

invention enables selecting out various compounds (which react with the LY6H receptor, inclusive of low molecular compounds, high molecular compounds, proteins, protein fragments, antigens, antibodies, etc.). Preferably, the
5 LY6H receptor as a whole is used. The LY6H receptor polypeptide or fragment thereof for use in such screening may have been immobilized on a solid matrix or be a free substance in a solution to be transported to the cell surface.

10 An example of the above pharmacoscreening is a screening system in which prokaryotic or eukaryotic host cells transformed stably with a recombinant DNA coding for an LY6H polypeptide, or a fragment thereof, are used in, preferably, a competitive binding assay. As an
15 alternative, said host cells, whether in the free form or as immobilized, are used in the standard binding assay. More particularly, the above pharmacoscreening may comprise reacting the LY6H receptor polypeptide, or a fragment thereof, with the LY6H polypeptide, or a
20 fragment thereof, in the presence of a candidate drug, to cause formation of a complex and detecting the degree of inhibition of the complex formation by the above candidate drug.

Thus, in accordance with the invention, there can
25 be provided a method for pharmacoscreening which

comprises contacting a candidate drug with the LY6H receptor polypeptide, or a fragment thereof and, then, detecting the presence of the resulting complex or the presence of a complex of the LY6H receptor polypeptide, or a fragment thereof, with a ligand by a per se known technique. Furthermore, by assaying LY6H receptor activity, it is possible to evaluate whether a candidate drug is capable of antagonizing the LY6H receptor and accordingly may modify the above-defined LY6H activity, i.e. may be able to modulate growth of neurons, or modulate protein-protein conjugation or complex-forming activity. In such a competitive binding assay, the LY6H receptor polypeptide, or a fragment thereof, is labeled. When the free LY6H receptor polypeptide or fragment thereof is separated from the protein-protein complex and the labeling amount of the free (non-complex-forming) substance is measured, the measured value serves as a yardstick of the binding of the test factor to the LY6H receptor. The measured value serves also as a measure of inhibition of the binding of the LY6H receptor to the LY6H polypeptide. By analyzing a small peptide (pseudopeptide) of the LY6H polypeptide in this manner, the candidate drug can be assayed as a substance having LY6H receptor antagonizing activity.

Another protocol for pharmacoscreening in

accordance with the invention is that of screening for a compound having an adequate binding affinity for the LY6H receptor polypeptide. Briefly, this procedure comprises synthesizing a large number of different test peptide
5 compounds on a solid support such as the surface of a plastic pin or other material, reacting the test peptide compounds with the LY6H receptor polypeptide and, after washing, detecting the binding reaction products of LY6H receptor polypeptide by a known method [e.g. PCT patent
10 publication No. WO 84-03564]. The purified LY6H receptor can be directly coated on the plate to be used in said pharmacoscreening procedure. The antibody may be captured with a non-neutralizing antibody against the polypeptide and the LY6H receptor polypeptide be immobilized on a
15 solid phase.

The invention is further directed to the use of a competitive pharmacoscreening assay. For the binding to the LY6H receptor polypeptide, or a fragment thereof, a neutralizing antibody capable of specific binding to the
20 LY6H receptor polypeptide is caused to compete with the candidate compound. By such a competitive reaction with the neutralizing antibody, the presence of any peptide having one or more antigenic determinants of the LY6H receptor polypeptide can be detected.

25 As a further method for drug screening, the LY6H

polypeptide of the invention or the LY6H gene product of the invention can be used in the screening for compounds which activate (agonists) or inhibit (antagonists or inhibitors) the activity of the LY6H polypeptide or LY6H
5 gene product.

By using the LY6H polypeptide or LY6H gene product of the invention, agonists or antagonists can be identified from cells, cell-free preparations, chemical libraries and naturally-occurring compositions. These
10 agonists or antagonists may be natural or modified substrates, ligands, enzymes or receptors of the LY6H polypeptide of the invention or structural or functional copies of the polypeptide of the invention [Coligan et al., Current Protocols in Immunology, 1(2), Chapter 5
15 (1991)].

In situ hybridization studies revealed the expression of LY6H gene of the invention in various tissues of the human normal brain, at particularly high levels in the hippocampus and entorhinal cortex which are
20 usually severely impaired in Alzheimer patients, and its expression level has been found to be considerably depressed in the temporal lobe inclusive of the hippocampus and entorhinal cortex of patients with Alzheimer's disease. It is, therefore, very likely that
25 this gene is associated with the onset and progression of

said disease.

Therefore, an agonist or antagonist of this LY6H protein or an LY6H gene product is expected to find application as a therapeutic or prophylactic drug for
5 neurodegenerative diseases such as Alzheimer's disease, Alzheimer type dementia, brain ischemia and Parkinson's disease.

Compounds obtainable by the screening for candidate drugs for said LY6H gene-related diseases have the
10 functions of the protein of the invention (the expression product of the gene of the invention), such as neuronal survival-supporting action, nerve elongating action, nerve regenerating action, neuroglia-activating action, etc. in the central and other nerve systems and brain
15 mnemonic (memory-forming) action, among other physiological actions, and, therefore, can be used as a therapeutic or prophylactic drug for various neurodegenerative diseases such as Alzheimer's disease, Alzheimer type dementia, brain ischemia and Parkinson's
20 disease. Thus, the proteins of the invention (inclusive of the gene expression products, partial peptides thereof, and salts thereof) are of use as reagents for the screening for compounds which promote the functions of the protein of the invention.

25 The invention provides a method of screening for

compounds which promote the functions of the protein of the invention (hereinafter each referred to sometimes as a functional enhancer of the protein of the invention). More particularly, the invention provides (a) a method of
5 screening for a functional enhancer of the protein of the invention which comprises contacting (1) the protein of the invention with nerve cells or a nerve tissue on one hand and (2) the protein and a test compound with said nerve cells or tissue on the other hand and comparing the
10 results and (b) a method of screening for a functional enhancer of the protein of the invention which comprises administering (1) the protein of the invention to a vertebrate on one hand and (2) the protein of the invention and a test compound to the vertebrate on the
15 other hand and comparing the results.

More particularly, in the above screening method (a), a physiological activity in the central or other nervous systems, such as neuronal survival-supporting activity, nerve elongating activity, nerve regenerating
20 activity or neuroglia-activating activity, is measured under the above conditions (1) and (2) and the results are compared. In the screening method (b), the mnemonic (memory-forming) activity in the brain, for instance, is measured under said two conditions (1) and (2) and the
25 results are compared.

The nerve cells (neurous and neuroglia) for use in the above screening include neuroblastoma cells, glioma cells, and their hybridoma cells (e.g. N18TG-2, IMR-32, GOTO (e.g. GOTO-P3), NB1, C6BU-1, U251, KNS42, KNS81 and NG108-15 cells, and PC12 cells having a potency of differentiation to nerve cells). The nerve tissue which can be used includes the mouse neuroepithelial cell, rat hippocampus primary culture cell, fetal mouse culture Prukinje cell, and mouse dorsal root ganglia. The test compound includes peptides, proteins, nonpeptide compounds, synthetic compounds, fermentation products, cell extracts, plant extracts, animal tissue extracts, and plasma. These compounds may be novel compounds or known compounds.

In carrying out said screening method (a), the protein of the invention (inclusive of a partial peptide thereof or a salt thereof) is dissolved or suspended in a screening buffer to prepare a sample of the protein of the invention. The buffer may be any buffer solution that does not interfere with the contact between the protein of the invention and the nerve cell or tissue (e.g. phosphate buffer, Tris-HCl buffer, etc. at pH about 4-10, preferably pH about 6-8). The duration of contact is usually about 1-10 days, preferably about 7-10 days. The contact temperature is usually about 37°C. The activities

of the protein of the invention in the central or other nervous systems, such as neuronal survival-supporting activity, nerve elongating activity, nerve regenerating activity, and neuroglia-activating activity, can be
5 determined by the routine methods such as visual assessment of axonal elongation, measurement of intracellular Ca^{2+} concentration, and the like.

Any test compound promoting any of said physiological activities, such as neuronal survival-
10 supporting activity, nerve elongating activity, nerve regenerating activity, neuroglia-activating activity, by at least about 20%, preferably not less than about 30%, more preferably not less than about 50%, still more preferably not less than about 70%, under the above-
15 mentioned condition (2) as compared with the condition (1) can be selected as a functional enhancer of the protein of the invention.

In carrying out the above screening method (b), the protein of the invention, alone or in combination with
20 the test compound, is administered to test animals by intravenous, subcutaneous or intramuscular injection or orally. The dosage of the protein of the invention for oral administration is generally about 0.1-100 mg/day, preferably about 1.0-50 mg/day, more preferably about
25 1.0-20 mg/day, per mammal (based on 50 kg body weight).

The parenteral dose should be selected according to the recipient and the method of administration but it is preferable to administer about 0.01-30 mg/day, preferably about 0.1-20 mg/day, more preferably about 0.1-10 mg/day, per mammal (50 kg body weight) by the intravenous route.

Test animals include such mammals as man, monkey, chimpanzee, mouse, rat, rabbit, sheep, swine, bovine, horse, cat and dog and fish (e.g. carp, salmon, herring, rainbow trout, goldfish, etc.).

The mnemonic (memory-forming) activity of the protein of the invention in the brain can be assayed in accordance with, for example, a water maze test protocol [Morris, R. G. M., J. Neurosci. Meth., 11, 47-60 (1984)]. Any test compound promoting the above mnemonic effect by not less than about 20%, preferably not less than 50%, more preferably not less than 70%, under said condition (2) as compared with said condition (1) is of use as a functional enhancer of the protein of the invention.

The screening kit as a further embodiment of the invention contains the protein of the invention (inclusive of the expression product of the gene, a partial peptide thereof, and any salt of either of them) as an essential component. A kit consists of the following components 1-4 is an example of the screening kit of the invention.

Component 1: Hanks solution as assay buffer

Component 2: Protein standard (protein of the invention or a salt thereof)

Component 3: Nerve cells or a nerve tissue (a culture of
5 said nerve cells or nerve tissue in a 24-well plate, 10^4
cell/well, as grown using Eagle's MEM, Hanks solution
under 5% CO₂ at 37°C)

Component 4: An inverted microscope for observation

The screening with the above screening kit can be
10 carried out as follows.

[Method]

The number per field of vision of axonal
elongation-positive cells in the well containing the test
compound is counted and compared with the number of
15 axonal elongation-positive cells in the control (test
compound-free) well and the difference is statistically
tested.

The compound or salt obtained by the screening
method or with the screening kit in accordance with the
20 invention is a member selected from the above-mentioned
class consisting of peptides, proteins, nonpeptide
compounds, synthetic compounds, fermentation products,
cell extracts, plant extracts, animal tissue extracts,
etc. and is a compound capable of promoting the function
25 of the protein of the invention. The compound that

promotes the functions of the protein of the invention as such may show physiological activities such as neuronal survival-supporting activity, nerve elongating activity, nerve regenerating activity, neuroglia-activating

5 activity, etc. and thereby promote the function of the protein of the invention or the like additively or synergistically or, although not showing such physiological activities by itself, may promote the function of the protein of the invention. Examples of the
10 salts of the compound include salts with physiologically acceptable bases (e.g. alkali metals) or acids (e.g. organic acids, inorganic acids). Particularly preferred are physiologically acceptable acid addition salts, such as salts with inorganic acids (e.g. hydrochloric acid,
15 phosphoric acid, hydrobromic acid, sulfuric acid) or organic acids (e.g. acetic acid, formic acid, propionic acid, fumaric acid, maleic acid, succinic acid, tartaric acid, citric acid, malic acid, oxalic acid, benzoic acid, methanesulfonic acid, benzenesulfonic acid).

20 The compound or salt which promotes the function of the protein of the invention is of value as a safe, low-toxicity therapeutic-prophylactic drug for various neurodegenerative diseases such as Alzheimer's disease, Alzheimer type dementia, brain ischemia and Parkinson's
25 disease.

The above screening procedure involves the use of cells which express the LY6H polypeptide on the cell surface or respond to the protein of the invention. Among such cells are cells derived from mammalian animals, yeasts, Drosophila and E. coli. The cells which express the LY6H polypeptide (or the cell membrane having the expressed polypeptide) or respond to the LY6H polypeptide is contacted with the test compound to observe the stimulation or inhibition of binding or functional response. Then, LY6H activity of cells contacted with the candidate compound is compared with that of similar cells not contacted.

The above assay can be carried out by detecting adhesion to cells harboring the LY6H polypeptide using a label directly or indirectly coupled to a candidate compound or in an assay system utilizing a competition with a label-competitive substance. In this manner, the binding of the candidate compound can be easily tested. Furthermore, using a detection system suited to cells bearing the LY6H polypeptide in such assays, it may be tested whether the candidate compound will produce a signal ascribable to activation of the LY6H polypeptide. The activation inhibitor is generally assayed in the presence of a known agonist and the effect of the candidate compound on the activation due to the agonist

is observed. The assay may comprise a simple procedure comprising mixing the candidate compound with a solution containing the LY6H polypeptide to form a mixture, determining the LY6H activity in the mixture, and
5 comparing the LY6H activity of the mixture with a standard.

The low molecular compound (agonist or antagonist) which binds to the LY6H protein can be obtained by a screening with BIACORE 2000, for instance [Markgren, P.
10 O., et al., Analytical Biochemistry, 265, 340-350 (1998)].

In accordance with the invention, for the purpose of developing a more active or stabilized LY6H polypeptide derivative or a drug which enhances or blocks the function of the LY6H polypeptide in vivo, it is
15 possible to construct a biologically active polypeptide or a structural analog thereof for interaction, such as an LY6H agonist, LY6H antagonist, LY6H inhibitor or the like. The structural analog mentioned above can be obtained, for example, by determining the three-
20 dimensional structure of a complex of LY6H polypeptide with another protein by X-ray crystallography, computer modeling or a combination of such techniques. Information on the structure of a structural analog can also be acquired by polypeptide modeling based on the structures
25 of homologous proteins.

To obtain said more active or stabilized LY6H polypeptide derivative, analysis by alanine scan can be employed. This method comprises substituting Ala for each amino acid residue to assess the influence of

5 substitution on peptide activity. Thus, as each amino acid residue of a peptide is thus analyzed, the region of importance to the activity or stability of the peptide is determined. By this method, it is possible to design a more active or stable LY6H polypeptide derivative.

10 It is also possible to isolate the target-specific antibody selected by the functional assay and analyze its crystal structure. As a rule, by this approach, the pharmacore providing a basis for subsequent drug design is obtained. By producing an anti-ideotypic antibody to
15 the functional pharmacologically active antibody, it is possible to identify and isolate a peptide from a chemically or biologically generated peptide bank. Therefore, it is predictable that the selected peptide may also serve as a pharmacore.

20 In this manner, it is possible to design and develop drugs having improved or stabilized LY6H activity or acting as inhibitors, agonists or antagonists of LY6H activity.

Evaluation of such a drug can be made by titrating
25 its effect on neuronal survival using primary culture

hippocampal neurons [Japan. J. Pharmacol, 53, 221-227
(1990)] or investigating its effect on neurodegenerative
lesions in Alzheimer model animals such as mutant β -
amyloid precursor protein gene or mutant presenilin 1
5 gene transgenic mice [Nature, 373, 523-527 (1995): Nature
Med., 5, 560-564 (1999)].

The compound thus obtained can be used not only as
a drug for Alzheimer's disease but also as a therapeutic
drug for cerebral infarction and other neurodegenerative
10 diseases.

Furthermore, in accordance with the invention, by
constructing LY6H gene-bearing knockout mice (transgenic
mice with LY6H knockout backgrounds), it is possible to
ascertain which site or sites of the nucleotide sequence
15 of the LY6H gene have influences on said multiple LY6H
activities in vivo, that is to say what functions the
expression products of LY6H gene and of a modified LY6H
gene have in vivo.

This method is a technique to intentionally modify
20 the genetic information of a living thing by utilizing
homologous recombinant genes, and includes a method using
mouse embryonic stem cells (ES cells) as an example
[Capecchi, M. R., Science, 244, 1288-1292 (1989)].

The method of constructing said mutant mice is by
25 now a routine technology for those skilled in the art,

and mutant mice can be easily constructed by applying a human wild-type LY6H gene or a mutant LY6H gene to a modified version of the above technology [Noda, Testuo (ed.): Jikken Igaku (Experimental Medicine), Supplement, 5 14(20) (1996), Yōdoshā]. Therefore, by utilizing this technique, it is possible to design and develop drugs having improved or stabilized LY6H activity or inhibitors, agonists, and antagonists of LY6H activity.

BRIEF DESCRIPTION OF THE DRAWING

10 Fig. 1 is a diagrammatic representation of Northern blots showing the pattern of expression of LY6H gene in various sites of the brain of a patient with Alzheimer's disease.

BEST MODE FOR CARRYING OUT THE INVENTION

15 The following examples are intended to illustrate the invention in further detail.

Example 1

(1) Cloning and DNA sequencing of human LY6H gene

The mRNA extracted from the human fetal brain was purchased from CLONTECH Laboratories and used as the 20 starting material. From this mRNA, a cDNA was synthesized and ligated into the vector λZAPII (Stratagene) to construct a cDNA library (Otsuka GEN Research Institute, Otsuka Pharmaceutical Co.). Using the in vivo excision 25 method [in vivo excision: Short, J. M., et al., Nucleic

Acids Res., 16, 7583-7600 (1988)], colonies of Escherichia coli bearing the human gene were formed on agar medium and randomly picked up to register the human gene-bearing E. coli clones in a 96-well microplate.

5 These clones were stored at -80°C.

Then, each registered clone was cultured in 1.5 ml of LB medium overnight and the DNA was extracted and purified using an automatic plasmid extractor PI-100 (Kurabo). The contaminated E. coli RNA was decomposed
10 with RNase and removed. Finally, 30 µl of a DNA solution was prepared and using a 2 µl portion, the approximate DNA size and amount were checked by the minigel method. A 7 µl portion was used for a sequencing reaction and the remaining 21 µl was stored as plasmid DNA at 4°C. By this
15 method, a cosmid which can also be used as a probe for FISH (fluorescence in situ hybridization) described below can be extracted by a minor modification of the program.

Then, a dideoxy terminator reaction of Sanger et al. using T3, T7 or a synthetic oligonucleotide primer
20 [Sanger, F., et al., Proc. Natl. Acad. Sci., USA., 74, 5463-5467 (1977)] or a cycle sequencing reaction [Carothers, A. M., et al., Bio. Techniques, 7, 494-499 (1989)] which is the dideoxy terminator reaction plus PCR was carried out. These are techniques for chain extension
25 with termination specific to 4 kinds of bases using a

small amount (about 0.1-0.5 µg) of plasmid DNA as the template.

Using an FITC (fluorescein isothiocyanate)-labeled primer as the sequence primer, about 25 cycles of reaction using Taq polymerase were carried out. Of the fluorescence-labeled DNA fragment, the sequence of about 400 nucleotides from the 5'-end of the cDNA was determined with the automatic DNA sequencer ALF™ DNA Sequencer (Pharmacia).

10 The 3'-nontranslated region is high in heterogeneity among genes and suited for differentiation of individual genes. Therefore, sequencing of the 3'-end region was also performed in some cases.

15 The huge nucleotide sequence information generated with the DNA sequencer was transmitted to the 64-bit computer DEC3400 for computerized homology analysis. This homology analysis was carried out by a database (GenBank, EMBL) search according to UWGCG's FASTA Program [Pearson, W. R. and Lipman, D. J., Proc. Natl. Acad. Sci., USA., 85,
20 2444-2448 (1988)].

Fujiwara et al. describe in detail about the above method of analysis for a human fetal brain cDNA library [Fujiwara, T., et al., DNA Res., 2, 107-111 (1991)].

25 The ESTs (expressed sequence tags: partial DNA sequences of the expressed gene fragment) randomly

selected from the human fetal brain cDNA library constructed as above were then sequenced.

The clone designated GEN-425D01 in the GenBank/EMBL sequence search according to the FASTA Program was found
5 to be highly homologous to the gene coding for the mouse Ly6 family protein.

Using a double-stranded DNA inserted into a vector (pBluescript vector; Stratagene) as a template and a
10 synthetic oligonucleotide as a primer, the nucleotide sequence of the cDNA inclusive of the whole coding region of the above clone was determined by Sanger's dideoxy chain termination method.

Sequencing with ABIPRISMTM377 automatic DNA
sequencer revealed that the cDNA sequence of the clone
15 obtained above contained a deduced amino acid coding region of 420 bases and the amino acid sequence encoded thereby had 140 amino acid residues. The nucleic acid sequence of the full-length cDNA clone was composed of 854 nucleotides. The full sequence is shown in SEQ ID
20 NO:3; the nucleotide sequence of the open reading frame is shown in SEQ ID NO:2; and the deduced amino acid sequence encoded by said nucleotide sequence is shown in SEQ ID NO:1.

The amino acid sequence of the human LY6H protein
25 was compared with the sequences of other Ly6 family

proteins, and the nucleotide sequence conserved in the amino acid translation initiation region [Kozak, M., J. Biol. Chem., 266, 19867-19870 (1991)] was compared with the 5'-region of the human LY6H gene. The initiation
5 codon thus determined was located in the position 99-101, which is the second ATG triplet, of the nucleotide sequence shown in SEQ ID NO:3. Moreover, the polyadenylation signal (AATAAA) was located in the position 832-837 of the same nucleotide sequence.

10 (2) Northern blot analysis

To define the expression profile of LY6H in tissues, a Northern blot analysis was performed using various human tissues.

In the Northern blot analysis, Human MTN (Multiple-
15 Tissue Northern) Blot I and II (CLONTECH) were used.

The cDNA fragment was amplified by PCR using a primer set of T3 and T7 promoter sequences.

The PCR amplification product of said GEN-425D01cDNA clone was labeled with [³²P]-dCTP (Random
20 Primed DNA Labeling Kit, Boehringer Mannheim GmbH) for use as a probe.

The blot containing the amplification product was prehybridized (under conditions according to the product protocol) and, then, subjected to hybridization according
25 to the product protocol.

The hybridization was performed at 65°C overnight in a solution composed of 1 M NaCl/50 mM Tris-HCl (pH 7.5)/2 × Denhardt's solution/10% dextran sulfate/1% SDS solution (containing 100 µg/ml denatured salmon sperm DNA). After wash twice with 2 × SSC/0.1% SDS at room temperature, the product was washed once with 0.1 × SSC/0.1% SDS at 65°C for 40 minutes. The filter was exposed against X-ray film (Kodak) at -70°C for 18 hours.

The above test was performed using the following adult human tissues: brain, pancreas, testis, small intestine, colon, thymus, prostate, ovary, heart, placenta, lung, liver, skeletal muscle, kidney, spleen, testis and peripheral blood leukocyte. As a result, transcripts of about 1 kb showing homology to LY6H were observed in the brain, pancreas, testis, small intestine, colon, thymus, prostate and ovary, particularly high in the brain.

(3) Localization of the gene on chromosome by FISH using cosmid clones

FISH for chromosomal localization was carried out using 0.5 µg of each cosmid DNA as a probe in accordance with the known method [Takahashi, E. et al., Hum. Genet., 86, 14-16 (1990)]. It was caught FISH signals by Provia 100 film (Fuji, ISO 100) or CCD Camera System (Applied Imaging Cyto Vision).

As a result, the human LY6H gene was found to be located on q24.3 of chromosome 8. Thus, GEN-425D01 was mapped on the chromosome band 8q24.3.

The antibodies against proteins belonging to the Ly6 family have been utilized in the purification of blood stem cells as a target of gene therapy [van de Rijn, M., et al., Proc. Natl. Acad. Sci., USA., 86, 4634-4638 (1989)], studies on the differentiation of blood cells [van de Rijn. M., et al., Proc. Natl. Acad. Sci., USA., 86, 4634-4638 (1989); Classon, B. J. and Coverdale, L., Proc. Natl. Acad. Sci., USA., 91, 5296-5300 (1994)], activation of immune cells [Malek, T. R., et al., J. Exp. Med., 164, 709-722 (1986)], inhibition of production of active immune cells [Haque, A., et al., Immunology, 69, 558-563 (1990)], and the like, and have also been found to have antitumor effects [Lu, L., et al., J. Immunol., 142, 719-725 (1989)]. The human LY6H gene provided in the present example enables detection of the expression of the gene in various tissues, production of the human LY6H protein by genetic engineering techniques, and construction of an antibody by utilizing the gene, hence enabling said purification of blood stem cells, research into the differentiation of blood cells, activation of immune cells, inhibition of activation of immune cells, and therapy of tumors.

Furthermore, the LY6H expressed at a high level in the brain enables a research into the differentiation of nerve cells, activation of neurons, and therapy of neural and mental diseases.

5 Screening for compounds with the human LY6H protein as the target is also made possible and the compounds thus obtained are as useful as the anti-human LY6H protein antibody.

Example 2

10 (1) Northern blot analysis in the brain tissues of a patient with Alzheimer's disease

Northern blot analysis was performed in accordance with Example 1 (2).

15 To investigate the expression of the LY6H gene in the brain tissues of patients with Alzheimer's disease, Northern blot analysis was made using the brain tissues of an Alzheimer patient and normal human brain tissues.

20 Northern blotting was performed using the human normal brain blot II and human Alzheimer blot II (both Invitrogen) and the LY6H gene expression in the various brain tissues, namely the frontal lobe, temporal lobe, parietal lobe, occipital lobe, pons, thalamus and corpus callosum, was compared between normal and Alzheimer brains.

25 The results are shown in Fig. 1.

As was pointed out in Example 1, the LY6H gene is expressed at a high level in the brain. The above analysis revealed that, while the expression of the gene was confirmed in various tissues of the human normal
5 brain, the gene was expressed at particularly high levels in the temporal lobe inclusive of the hippocampus and entorhinal cortex which are known to be impaired severely in patients with Alzheimer's disease while marked decreases were found in the temporal lobe inclusive of
10 the hippocampus and entorhinal cortex in the patient with Alzheimer's disease, indicating that it is very likely that the gene is involved in the onset and progression of this disease.

Therefore, the LY6H gene sense strand, LY6H
15 expression product, and LY6H protein are expected to find application as therapeutic drugs for Alzheimer's disease, Alzheimer type dementia, brain ischemia and Parkinson's disease.

Furthermore, agonists and antagonists of LY6H
20 protein are also expected to be of use as therapeutic drugs for Alzheimer's disease and other diseases.

Example 3

(1) Construction of an LY6H expression vector

The LY6H cDNA obtained by in vivo excision method
25 is cleaved with MvII and XhoI to give an about 800-base

fragment. This fragment, containing the entire coding region of the LY6H gene shown in SEQ ID NO:1, is ligated to the EcoRV/XhoI-cleaved pAc5.1/V5-HisA (Invitrogen) to construct an expression vector (pAC/LY6H expression vector).

(2) Expression and purification of the active ingredient protein of the invention

The pAC/LY6H expression vector DNA and pCoHYGRO vector (Invitrogen) DNA are admixed in a ratio of 19:1 and introduced into fruit fly (Schneider 2) cells by calcium phosphate transfection. After the cells are cultured in 10% fetal calf serum-DES expression medium (Invitrogen) at 23°C for 48 hours, 300 µg/ml of hygromycin (Hygromycini B, Boehringer Mannheim) is added to the culture and the selection of drug-resistant cell clones is performed for 2 weeks. A stable transformant is subjected to stationary culture at a concentration of 5×10^6 cells/ml using 20 Falcon 5000 culture flasks (Becton Dickinson) containing 20 ml of 10% fetal calf serum-DES expression medium (Invitrogen) and the cultured cells are harvested. After washing twice with phosphate-buffered saline (PBS), the cells are suspended in PBS containing 2% bovine serum albumin and 0.5 U/ml of phosphatidyl-inositol-specific phospholipase C (PIPLC) and cultured at 37°C for 1 hour. From the supernatant of the culture, the

objective protein can be purified by ion exchange column chromatography or the like.

(3) Isolation and culture of hippocampal neurons

The whole brain is aseptically isolated from fetal
5 SD rats on embryonic day 18 and the hippocampus is
excised. The excised tissue is cut to thin slices with a
surgical knife and incubated for enzymatic treatment in
PBS containing 0.25% trypsin and 0.002% DNase I at 37°C
for 20 minutes. After the enzymatic reaction is stopped
10 by adding fetal calf serum, the aspiration-ejection of
the cell digest with a pipette having a plastic tip is
repeated 3 times to disperse the cells. The cell
dispersion is passed through a filter consisting of 2
stacked sheets of lens paper to remove the undigested
15 tissue and centrifuged at 1000 rpm for 5 minutes. The
cells are washed with DMEM (Gibco) and seeded on a poly-
L-lysine (Sigma)-coated 96-well plate containing 10% FCS-
DMEM at a final concentration of 2×10^5 cells/cm².

(4) Treatment with the active ingredient protein of the
20 invention

The above cells are cultured for 24 hours and after
the culture medium is changed to 1% N₂ Supplement (Gibco)-
DMEM, the active ingredient protein of the invention as
prepared under (2) is added (the invention group).

25 For comparison, the active ingredient protein of

the invention is heat-treated in a boiling water bath for 5 minutes and added (the boiled protein group).

(5) Evaluation of hippocampal neuronal survival

The cells (culture) in each group as prepared under
5 (4) are cultured for 72 hours. Then, the hippocampal neuronal survival-supporting effect of the active ingredient protein of the invention can be evaluated by the MTT [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide] assay. This MTT assay may be
10 performed using Promega's "CellTiter 96" Assay System, for instance.

(6) Isolation and culture of midbrain neurons

The whole brain is aseptically isolated from fetal SD rats on embryonic day 14 and the ventral midbrain is
15 excised. The tissue is cut to thin slices with a surgical knife and incubated for enzymatic treatment in phosphate-buffered saline (PBS) containing 0.25% trypsin and 0.002% DNase I at 37°C for 20 minutes. After the enzymatic reaction is stopped by adding fetal calf serum, the
20 aspiration-ejection of the cell digest with a pipette having a plastic tip is repeated 3 times to disperse the cells. The cell dispersion is passed through a filter consisting of 2 stacked sheets of lens paper to remove the undigested tissue and centrifuged at 1000 rpm for 5
25 minutes. The cells are washed with DMEM/F12 (Gibco) and

seeded on a poly-L-lysine-coated 96-well plate containing 10% FCS-DMEM/F12 at a final concentration of 3×10^5 cells/cm².

5 (7) Treatment with the active ingredient protein of the invention

 The cells prepared under (6) are cultured for 24 hours and after the culture medium is changed to 1% N₂ Supplement (Gibco)-DMEM/F12, the active ingredient protein of the invention as prepared under (2) is added
10 (the invention group).

 For comparison, the active ingredient protein of the invention is heat-treated in a boiling water bath for 5 minutes and added (the boiled protein group).

15 (8) Evaluation of midbrain neuronal survival supporting effect

 The cells (culture) in each group as prepared under (7) are cultured for 72 hours. Then, the midbrain neuronal survival-supporting effect of the active ingredient protein of the invention can be evaluated by
20 the MTT [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide] assay. This MTT assay may be performed using Promega's "CellTiter 96" Assay System, for instance.

25 (9) Evaluation of dopaminergic neuronal survival-supporting effect

The cells (culture) in each group as prepared under (7) are cultured for 72 hours and, then, fixed by allowing them to stand in 4% paraformaldehyde-PBS at room temperature for 15 minutes. Thereafter, using 1% Triton X100/PBS, it is passed through a membrane.

To prevent nonspecific binding of the antibody, the cells are incubated in 10% goat serum-PBS for 1 hour and, then, using an anti-tyrosine hydroxylase polyclonal antibody (Chemicon; diluted 1000-fold with PBS), the cells are incubated at 4°C for 16 hours. After the antibody solution is removed, the cells are washed with PBS and, with peroxidase-labeled dextran polymer-conjugated goat anti-rabbit immunoglobulin (Dako) added, the cells are incubated at room temperature for 1 hour.

The tyrosine hydroxylase-positive cells can be detected by the color reaction using diaminobenzidine as the substrate. Using the number of tyrosine hydroxylase-positive cells as a marker, the dopaminergic neuronal survival-supporting effect can be evaluated.

INDUSTRIAL APPLICABILITY

The present invention provides a novel brain-specific gene and a protein encoded thereby and, by utilizing them, technologies of value to the purification of blood stem cells, research into the differentiation of blood cells, activation of immune cells, inhibition of

production of active immune cells, and therapy of tumors
can be provided. Also, the present invention provides
novel genes having physiologic activities such as brain
neuronal survival-supporting activity, nerve elongating
5 activity, nerve regenerating activity, neuroglia-
activating activity and brain memory-forming activity.

In view of the marked depression of its expression
level in the temporal lobe of the brain of patients with
Alzheimer's disease, the gene of the invention is
10 considered to inhibit neurodegenerative changes of the
tissue, thus being of use as a gene therapy drug.
Moreover, the expression product of the gene of the
invention finds application as a prophylactic and
therapeutic drug for such neurodegenerative diseases.

CLAIMS

1. A gene comprising a nucleotide sequence coding for the following protein (a) or (b):

(a) a protein having the amino acid sequence shown in SEQ

5 ID NO:1

(b) a protein having an amino acid sequence derived from the amino acid sequence shown in SEQ ID NO:1 by deletion, substitution or addition of one or plurality amino acids and having at least one physiologic activity selected
10 from the group consisting of neuronal survival-supporting activity, nerve elongating activity, nerve regenerating activity, neuroglia-activating activity and brain memory-forming activity.

2. The gene according to Claim 1 wherein the
15 nucleotide sequence is shown in SEQ ID NO:2.

3. A gene comprising the following polynucleotide

(a) or (b):

(a) a polynucleotide containing the nucleotide sequence shown in SEQ ID NO:3

20 (b) a polynucleotide which hybridizes under stringent conditions with a DNA having the nucleotide sequence shown in SEQ ID NO:3.

4. The gene according to any one of Claims 1-3 wherein the gene is a human gene.

25 5. A gene expression vector comprising the gene

according to claim 2 or 3.

6. A host cell comprising the gene expression vector according to Claim 5.

7. An expression product which is expressed by the
5 host cell according to Claim 6.

8. A protein which is encoded by the gene according to Claim 1.

9. An antibody having a binding affinity for the expression product according to Claim 7 or the protein
10 according to Claim 8.

10. An expression product, which comprises the whole or part of a gene of the nucleotide sequence shown in SEQ ID NO:2 and having at least one physiological activity selected from the group consisting of neuronal
15 survival-supporting activity, nerve elongating activity, nerve regenerating activity, neuroglia-activating activity and brain memory-forming activity.

11. A therapeutic and prophylactic composition for neurodegenerative disease, which comprises the protein
20 according to Claim 8, an equivalent protein having a partial amino acid sequence thereof, or the expression product according to Claim 10 as an active ingredient in combination with a pharmaceutical carrier.

12. The therapeutic and prophylactic composition
25 for neurodegenerative disease according to Claim 11

wherein the neurodegenerative disease is selected from the group consisting of Alzheimer's disease, Alzheimer type dementia, brain ischemia and Parkinson's disease.

13. A sense strand oligonucleotide comprising at
5 least 20 consecutive constituent nucleotides of the nucleotide sequence shown in SEQ ID NO:2.

14. A gene therapy composition comprising the sense strand oligonucleotide according to Claim 13 as an active ingredient in combination with a pharmaceutical carrier.

10 15. A gene-specific probe comprising an oligonucleotide sequence of at least 10 consecutive constituent nucleotides of the nucleotide sequence shown in SEQ ID NO:2.

16. A method of screening for candidate compounds,
15 which comprises using the protein according to Claim 8, an equivalent protein having a partial amino acid sequence thereof, or the expression product according to Claim 10, said candidate compounds bind to or influence the protein, equivalent protein or gene product.

20 17. A method for therapy and prophylaxis of neurodegenerative disease, which comprises administering to a patient an effective amount of the protein according to Claim 8, an equivalent protein having a partial amino acid sequence thereof, or the expression product
25 according to Claim 10.

18. The method for therapy and prophylaxis of neurodegenerative disease according to Claim 17 wherein the neurodegenerative disease is selected from the group consisting of Alzheimer's disease, Alzheimer type
5 dementia, brain ischemia and Parkinson's disease.


19. A method of gene therapy, which comprises administering to a patient an effective amount of the sense strand oligonucleotide according to Claim 13.

ABSTRACT

The invention provides a brain-specific gene useful
in treating Alzheimer's disease, for instance, which
comprises a nucleotide sequence coding for the amino
5 acid sequence shown in SQ ID NO:1 and fragments thereof;
an expression vector comprising the gene; a host cell
comprising the expression vector; an expression product
of the gene; an antibody against the product; a
therapeutic and prophylactic composition for
10 neurodegenerative disease; and the like.

1/1

F I G. 1



Declaration and Power of Attorney for Patent Application

特許出願宣言書及び委任状

Japanese Language Declaration

日本語宣言書

下記の氏名の発明者として、私は以下の通り宣言します。

As a below named inventor, I hereby declare that:

私の住所、私書箱、国籍は下記の私の氏名の後に記載された通りです。

My residence, post office address and citizenship are as stated next to my name,

下記の名称の発明に関して請求範囲に記載され、特許出願している発明内容について、私が最初かつ唯一の発明者(下記の氏名が一つの場合)もしくは最初かつ共同発明者であると(下記の名称が複数の場合)信じています。

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

LY6H GENE

上記発明の明細書(下記の欄でX印がついていない場合は、本書に添付)は、

the specification of which is attached hereto unless the following box is checked:

☐ ____月 ____日に提出され、米国出願番号または特許協定条約

☒ was filed on September 16, 1999
as United States Application Number or
PCT International Application Number

国際出願番号を _____ とし、

PCT/JP99/05039 and was amended on
_____ (if applicable).

(該当する場合) _____ に訂正されました。

私は、特許請求範囲を含む上記訂正後の明細書を検討し、内容を理解していることをここに表明します。

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

私は、連邦規則法典第37編第1条56項に定義されるとおり、特許資格の有無について重要な情報を開示する義務があることを認めます。

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56.

Japanese Language Declaration

(日本語宣言書)

私は、米国法典第35編第119条(a)-(d)項又は第365条(b)項に基づき下記の、米国以外の国の少なくとも一カ国を指定している特許協力条約第365条(a)項に基づく国際出願、又は外国での特許出願もしくは発明者証の出願についての外国優先権をここに主張するとともに、優先権を主張している本出願の前に出願された特許または発明者証の外国出願を以下に、枠内をマークすることで、示しています。

I hereby claim foreign priority under Title 35, United States Code, Section 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed.

Prior Foreign Applications

外国での先行出願

Priority Not Claimed

優先権主張なし

1998-263550
(Number)
(番号)

Japan
(Country)
(国名)

17/09/1998
(Day/Month/Year Filed)
(出願年月日)

☐

(Number)
(番号)

(Country)
(国名)

(Day/Month/Year Filed)
(出願年月日)

☐

(Number)
(番号)

(Country)
(国名)

(Day/Month/Year Filed)
(出願年月日)

☐

私は、第35編米国法典119条(e)項に基づいて下記の米国特許出願規定に記載された権利をここに主張致します。

I hereby claim the benefit under Title 35, United States Code, Section 119(e) of any United States provisional application(s) listed below.

(Application No.)
(出願番号)

(Filing Date)
(出願日)

(Application No.)
(出願番号)

(Filing Date)
(出願日)

私は、下記の米国法典第35編第120条に基づいて下記の米国特許出願に記載された権利、又は米国を指定している特許協力条約第365条(c)に基づく権利をここに主張します。又、本出願の各請求範囲の内容が米国法典第35編第112条第1項又は特許協力条約で規定された方法で先行する米国特許出願に開示されていない限り、その先行米国出願書提出日以降で本出願書の日本国内又は特許協力条約国際出願提出日までの期間中に入手された、連邦規則法典第37編第1条第56項で定義された特許資格の有無に関する重要な情報について開示義務があることを認識しています。

I hereby claim the benefit of Title 35, United States Code Section 120 of any United States application(s), or 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of Title 35, United States Code Section 112, I acknowledge the duty to disclose any material information which is material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application:

(Application No.)
(出願番号)

(Filing Date)
(出願日)

(Status: Patented, Pending, Abandoned)
(現況: 特許許可済、係属中、放棄済)

(Application No.)
(出願番号)

(Filing Date)
(出願日)

(Status: Patented, Pending, Abandoned)
(現況: 特許許可済、係属中、放棄済)

私は、私自身の知識に基づいて本宣言中で私が行う表明が真実であり、かつ私の入手した情報と私の信ずるところに基づく表明が全て真実であると信じていること、さらに故意になされた虚偽の表明及びそれと同等の行為は米国法典第18編第1001条に基づき、罰金または拘禁、もしくはその両方により処罰されること、そしてそのような故意による虚偽の声明を行えば、出願した、又は既に許可された特許の有効性が失われることを認識し、よってここに上記のごとく宣誓を致します。

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Japanese Language Declaration

(日本語宣言書)

委任状: 私は、下記の発明者として、本出願に関する一切の
手続きを米国特許商標局に対して遂行する弁理士又は代理
人として、下記のことを指名致します。(弁理士、又は代理人の
氏名及び登録番号を明記のこと)

POWER OF ATTORNEY: As a named inventor, I hereby
appoint the following attorney(s) and/or agent(s) to
prosecute this application and transact all business in the
Patent and Trademark Office connected therewith (list
name and registration number)

John H. Mion, Reg. No. 18,879; Thomas J. Macpeak, Reg. No. 19,292; Robert J. Seas, Jr., Reg. No. 21,092;
Darryl Mexic, Reg. No. 23,063; Robert V. Sloan, Reg. No. 22,775; Peter D. Olexy, Reg. No. 24,513; J. Frank
Osha, Reg. No. 24,625; Waddell A. Biggart, Reg. No. 24,861; Louis Gubinsky, Reg. No. 24,835; Neil B.
Siegel, Reg. No. 25,200; David J. Cushing, Reg. No. 28,703; John R. Inge, Reg. No. 26,916; Joseph J. Ruch,
Jr., Reg. No. 26,577; Sheldon I. Landsman, Reg. No. 25,430; Richard C. Turner, Reg. No. 29,710; Howard L.
Bernstein, Reg. No. 25,665; Alan J. Kasper, Reg. No. 25,426; Kenneth J. Burchfiel, Reg. No. 31,333; Gordon
Kit, Reg. No. 30,764; Susan J. Mack, Reg. No. 30,951; Frank L. Bernstein, Reg. No. 31,484; Mark Boland,
Reg. No. 32,197; William H. Mandir, Reg. No. 32,156; Scott M. Daniels, Reg. No. 32,562; Brian W. Hannon,
Reg. No. 32,778; Abraham J. Rosner, Reg. No. 33,276; Bruce E. Kramer, Reg. No. 33,725; Paul F. Neils, Reg.
No. 33,102 and Brett S. Sylvester, Reg. No. 32,765

書類送付先:

Send Correspondence to:

SUGHRUE, MION, ZINN, MACPEAK & SEAS, PLLC
2100 Pennsylvania Avenue, N.W., Washington, D.C. 20037-3202

直通電話連絡先: (名称及び電話番号)

Direct Telephone Calls to: (name and telephone number)

(202)293-7060

唯一又は第一発明者名	Full name of sole or first inventor <u>MORIE Masato</u>		
発明者の署名	日付	Inventor's signature <u>Masato Morie</u>	Date Feb. 15, 2001
住所	Residence <u>Naruto-shi, Tokushima 772-0051, Japan</u>		
国籍	Citizenship Japan		
郵便の宛先	Post office address 261, Aza Minami, Takashima, Naruto-cho, Naruto-shi, Tokushima 772-0051, Japan		
第二共同発明者名(該当する場合)	Full name of second joint inventor, if any <u>OKUTOMI Keiichi</u>		
第二発明者の署名	日付	Second inventor's signature <u>Keiichi Okutomi</u>	Date Feb. 15, 2001
住所	Residence <u>Tokushima-shi, Tokushima 771-0128, Japan</u>		
国籍	Citizenship Japan		
郵便の宛先	Post office address Arupenroze 701, 51-1, Kamibetsukukita, Kawauchi-cho, Tokushima-shi, Tokushima 771-0128, Japan		

(第三以降の共同発明者についても同様に記載し、署名をするこ (Supply similar information and signature for third and
と。 subsequent joint inventors.)

Japanese Language Declaration (日本語宣言書)

第三共同発明者名 (該当する場合)		Full name of third joint inventor, if any TANIGUCHI Yoshihiro	
第三発明者の署名	日付	Third inventor's signature Yoshihiro Taniguchi	Date Feb. 15, 2001
住所		Residence Tokushima-shi, Tokushima 771-0125, Japan	
国籍		Citizenship Japan	
郵便の宛先		Post office address Rafore II 706, 5-1, Kanaoka, Kawauchi-cho, Tokushima-shi, Tokushima 771-0125, Japan	
第四共同発明者名 (該当する場合)		Full name of fourth joint inventor, if any SUZUKI Mikio	
第四発明者の署名	日付	Fourth inventor's signature Mikio Suzuki	Date Feb. 15, 2001
住所		Residence Tokushima-shi, Tokushima 771-0130, Japan	
国籍		Citizenship Japan	
郵便の宛先		Post office address Imagireryo, 463-30, Kagasuno, Kawauchi-cho, Tokushima-shi, Tokushima 771-0130, Japan	
第五共同発明者名 (該当する場合)		Full name of fifth joint inventor, if any OHBUCHI Yutaka	
第五発明者の署名	日付	Fifth inventor's signature Yutaka Ohbuchi	Date Feb. 15, 2001
住所		Residence Tokushima-shi, Tokushima 771-0130, Japan	
国籍		Citizenship Japan	
郵便の宛先		Post office address Imagireryo, 463-30, Kagasuno, Kawauchi-cho, Tokushima-shi, Tokushima 771-0130, Japan	
第六共同発明者名 (該当する場合)		Full name of sixth joint inventor, if any	
第六発明者の署名	日付	Sixth inventor's signature	Date
住所		Residence	
国籍		Citizenship	
郵便の宛先		Post office address	

09/787360

PCT
09RAW SEQUENCE LISTING
PATENT APPLICATION: US/09/787,360DATE: 03/28/2001
TIME: 10:54:03Input Set : A:\Q63396.txt
Output Set: N:\CRF3\03282001\I787360.raw

3 <110> APPLICANT: OTSUKA PHARMACEUTICAL CO., LTD.
 5 <120> TITLE OF INVENTION: LY6H GENE
 7 <130> FILE REFERENCE: Q63396
 C--> 9 <140> CURRENT APPLICATION NUMBER: US/09/787,360
 C--> 9 <141> CURRENT FILING DATE: 2001-03-16
 9 <150> PRIOR APPLICATION NUMBER: JP 1998-263550
 10 <151> PRIOR FILING DATE: 1998-09-17
 12 <160> NUMBER OF SEQ ID NOS: 3
 14 <170> SOFTWARE: PatentIn version 3.0
 16 <210> SEQ ID NO: 1
 17 <211> LENGTH: 140
 18 <212> TYPE: PRT
 19 <213> ORGANISM: Homo sapiens
 21 <220> FEATURE:
 22 <221> NAME/KEY: misc_feature
 23 <223> OTHER INFORMATION: human embryonic brain
 26 <400> SEQUENCE: 1
 28 Met Leu Pro Ala Ala Met Lys Gly Leu Gly Leu Ala Leu Leu Ala Val
 29 1 5 10 15
 31 Leu Leu Cys Ser Ala Pro Ala His Gly Leu Trp Cys Gln Asp Cys Thr
 32 20 25 30
 34 Leu Thr Thr Asn Ser Ser His Cys Thr Pro Lys Gln Cys Gln Pro Ser
 35 35 40 45
 37 Asp Thr Val Cys Ala Ser Val Arg Ile Thr Asp Pro Ser Ser Ser Arg
 38 50 55 60
 40 Lys Asp His Ser Val Asn Lys Met Cys Ala Ser Ser Cys Asp Phe Val
 41 65 70 75 80
 43 Lys Arg His Phe Phe Ser Asp Tyr Leu Met Gly Phe Ile Asn Ser Gly
 44 85 90 95
 46 Ile Leu Lys Val Asp Val Asp Cys Cys Glu Lys Asp Leu Cys Asn Gly
 47 100 105 110
 49 Ala Ala Gly Ala Gly His Ser Pro Trp Ala Leu Ala Gly Gly Leu Leu
 50 115 120 125
 52 Leu Ser Leu Gly Pro Ala Leu Leu Trp Ala Gly Pro
 53 130 135 140
 55 <210> SEQ ID NO: 2
 56 <211> LENGTH: 420
 57 <212> TYPE: DNA
 58 <213> ORGANISM: Homo sapiens
 60 <220> FEATURE:
 61 <221> NAME/KEY: misc_feature
 62 <223> OTHER INFORMATION: human embryonic brain
 65 <400> SEQUENCE: 2
 66 atgctgcctg cagccatgaa gggcctcggc ctggcgctgc tggcgcctct gctgtgctcg 60
 68 ggcgccgctc atggcctgtg gtgccaggac tgcaccctga ccaccaactc cagccattgc 120
 70 accccaagc agtgccagcc gtccgacacg gtgtgtgcca gtgtccgaat caccgatccc 180
 72 agcagcagca ggaaggatca ctcggtgaac aagatgtgtg cctcctcctg tgaattcggt 240

ENTERED

RAW SEQUENCE LISTING

DATE: 03/28/2001

PATENT APPLICATION: US/09/787,360

TIME: 10:54:03

Input Set A:\Q63396.txt

Output Set: N:\CRF3\03282001\I787360.raw

```

74 aagcgacact tttctcaga ctatctgatg gggtttatta actctgggat cttaaaggtc 300
76 gacgtggact gctgcgagaa ggatttgtgc aatggggcgg caggggcagg gcacagcccc 360
78 tgggccctgg ccggggggct cctgctcagc ctggggcctg cctcctcttg ggctgggccc 420
81 <210> SEQ ID NO: 3
82 <211> LENGTH: 854
83 <212> TYPE: DNA
84 <213> ORGANISM: Homo sapiens
86 <220> FEATURE:
87 <221> NAME/KEY: misc_feature
88 <223> OTHER INFORMATION: human embryonic brain
91 <220> FEATURE:
92 <221> NAME/KEY: exon
93 <222> LOCATION: (99)..(518)
95 <400> SEQUENCE: 3
96 acgcgcgccg agcccgaggc ggcgacaccc ccgggatgct tgcgccccag aggaccgcgc 60
98 cccaagccc ccgcgcgcc ccaggccca ccgggagc' atg ctg cct gca gcc atg 116
99 Met Leu Pro Ala Ala Met
100 1 5
102 aag ggc ctc ggc ctg gcg ctg ctg gcc gtc ctg ctg tgc tgc gcg ccc 164
103 Lys Gly Leu Gly Leu Ala Leu Leu Ala Val Leu Leu Cys Ser Ala Pro
104 10 15 20
106 gct cat ggc ctg tgg tgc cag gac tgc acc ctg acc acc aac tcc agc 212
107 Ala His Gly Leu Trp Cys Gln Asp Cys Thr Leu Thr Thr Asn Ser Ser
108 25 30 35
110 cat tgc acc cca aag cag tgc cag ccg tcc gac acg gtg tgt gcc agt 260
111 His Cys Thr Pro Lys Gln Cys Gln Pro Ser Asp Thr Val Cys Ala Ser
112 40 45 50
114 gtc cga atc acc gat ccc agc agc agc agg aag gat cac tgc gtg aac 308
115 Val Arg Ile Thr Asp Pro Ser Ser Ser Arg Lys Asp His Ser Val Asn
116 55 60 65 70
118 aag atg tgt gcc tcc tcc tgt gac ttc gtt aag cga cac ttt ttc tca 356
119 Lys Met Cys Ala Ser Ser Cys Asp Phe Val Lys Arg His Phe Phe Ser
120 75 80 85
122 gac tat ctg atg ggg ttt att aac tct ggg atc tta aag gtc gac gtg 404
123 Asp Tyr Leu Met Gly Phe Ile Asn Ser Gly Ile Leu Lys Val Asp Val
124 90 95 100
126 gac tgc tgc gag aag gat ttg tgc aat ggg gcg gca ggg gca ggg cac 452
127 Asp Cys Cys Glu Lys Asp Leu Cys Asn Gly Ala Ala Gly Ala Gly His
128 105 110 115
130 agc ccc tgg gcc ctg gcc ggg ggg ctc ctg ctc agc ctg ggg cct gcc 500
131 Ser Pro Trp Ala Leu Ala Gly Gly Leu Leu Leu Ser Leu Gly Pro Ala
132 120 125 130
134 ctc ctc tgg gct ggg ccc'tgatgtctcc tccttcccac ggggcttctg 548
135 Leu Leu Trp Ala Gly Pro
136 135 140
138 agcttgctcc cctgagcctg tggtgcct ctccccagcc tggcgtggct ggggctgggg 608
140 gcagccttgg ccagctccg tggctgtggc ctgtggctct cactcctccc ccgacgtgaa 668
142 gcctccctgt ctctccgcca gctctgagtc ccaggcagct ggacatctcc aggaaaccag 728
144 gccatctggg caggaggcct ggggatgagg gtgggggggg acccccaggt cccggagggg 788

```

RAW SEQUENCE LISTING

DATE: 03/28/2001

PATENT APPLICATION: US/09/787,360

TIME: 10:54:03

Input Set : A:\Q63396.txt

Output Set: N:\CRF3\03282001\I787360.raw

146 aagtgaagca acagcccagc tggaaggcg ttttctgcgg agaaataaag tcacttttga 848
148 gtcctg 854

146 aagtgaagca acagcccagc tggaaggcg ttttctgcgg agaaataaag tcacttttga 848
148 gtcctg 854

VERIFICATION SUMMARY

DATE: 03/28/2001

PATENT APPLICATION. US/09/787,360

TIME: 10:54:04

Input Set : A:\Q63396.txt

Output Set: N:\CRF3\03282001\I787360.raw

L:9 M:270 C: Current Application Number differs, Replaced Current Application No
L:9 M:271 C: Current Filing Date differs, Replaced Current Filing Date

03/28/2001 10:54:04

SEQUENCE LISTING

<110> OTSUKA PHARMACEUTICAL CO., LTD.

<120> LY6H GENE

<130> Q63396

<150> JP 1998-263550

<151> 1998-09-17

<160> 3

<170> PatentIn version 3.0

<210> 1

<211> 140

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> human embryonic brain

<400> 1

Met Leu Pro Ala Ala Met Lys Gly Leu Gly Leu Ala Leu Leu Ala Val
1 5 10 15

Leu Leu Cys Ser Ala Pro Ala His Gly Leu Trp Cys Gln Asp Cys Thr
20 25 30

Leu Thr Thr Asn Ser Ser His Cys Thr Pro Lys Gln Cys Gln Pro Ser
35 40 45

Asp Thr Val Cys Ala Ser Val Arg Ile Thr Asp Pro Ser Ser Ser Arg
50 55 60

Lys Asp His Ser Val Asn Lys Met Cys Ala Ser Ser Cys Asp Phe Val
65 70 75 80

Lys Arg His Phe Phe Ser Asp Tyr Leu Met Gly Phe Ile Asn Ser Gly
85 90 95

Ile Leu Lys Val Asp Val Asp Cys Cys Glu Lys Asp Leu Cys Asn Gly
100 105 110

Ala Ala Gly Ala Gly His Ser Pro Trp Ala Leu Ala Gly Gly Leu Leu
115 120 125

Leu Ser Leu Gly Pro Ala Leu Leu Trp Ala Gly Pro
130 135 140

<210> 2

<211> 420

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature
<223> human embryonic brain

<400> 2
atgctgcctg cagccatgaa gggcctcggc ctggcgctgc tggccgtcct gctgtgctcg 60
gcgcccgtc atggcctgtg gtgccaggac tgcaccctga ccaccaactc cagccattgc 120
accccaaagc agtgccagcc gtccgacacg gtgtgtgcc gtgtccgaat caccgatccc 180
agcagcagca ggaaggatca ctcggtgaac aagatgtgtg cctcctcctg tgacttcggt 240
aagcgacact ttttctcaga ctatctgatg gggtttatta actctgggat cttaaaggtc 300
gacgtggact gctgcgagaa ggatttgtgc aatggggcgg caggggcagg gcacagcccc 360
tggggcctgg ccgggggggt cctgctcagc ctggggcctg ccctcctctg ggctgggccc 420

<210> 3
<211> 854
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> human embryonic brain

<220>
<221> exon
<222> (99)..(518)

<400> 3
acgcgcgccg agcccgaggt gcggacaccc ccgggatgct tgcgccccag aggacccgcg 60
ccccaagccc ccgcgccgcc ccaggccca cccggagc atg ctg cct gca gcc atg 116
Met Leu Pro Ala Ala Met
1 5
aag ggc ctc ggc ctg gcg ctg ctg gcc gtc ctg ctg tgc tcg gcg ccc 164
Lys Gly Leu Gly Leu Ala Leu Leu Ala Val Leu Leu Cys Ser Ala Pro
10 15 20
gct cat ggc ctg tgg tgc cag gac tgc acc ctg acc acc aac tcc agc 212
Ala His Gly Leu Trp Cys Gln Asp Cys Thr Leu Thr Thr Asn Ser Ser
25 30 35
cat tgc acc cca aag cag tgc cag ccg tcc gac acg gtg tgt gcc agt 260
His Cys Thr Pro Lys Gln Cys Gln Pro Ser Asp Thr Val Cys Ala Ser
40 45 50
gtc cga atc acc gat ccc agc agc agc agg aag gat cac tcg gtg aac 308
Val Arg Ile Thr Asp Pro Ser Ser Ser Arg Lys Asp His Ser Val Asn
55 60 65 70
aag atg tgt gcc tcc tcc tgt gac ttc gtt aag cga cac ttt ttc tca 356
Lys Met Cys Ala Ser Ser Cys Asp Phe Val Lys Arg His Phe Phe Ser
75 80 85

gac tat ctg atg ggg ttt att aac tct ggg atc tta aag gtc gac gtg	404
Asp Tyr Leu Met Gly Phe Ile Asn Ser Gly Ile Leu Lys Val Asp Val	
90 95 100	
gac tgc tgc gag aag gat ttg tgc aat ggg gcg gca ggg gca ggg cac	452
Asp Cys Cys Glu Lys Asp Leu Cys Asn Gly Ala Ala Gly Ala Gly His	
105 110 115	
agc ccc tgg gcc ctg gcc ggg ggg ctc ctg ctc agc ctg ggg cct gcc	500
Ser Pro Trp Ala Leu Ala Gly Gly Leu Leu Leu Ser Leu Gly Pro Ala	
120 125 130	
ctc ctc tgg gct ggg ccc tgatgtctcc tccttcccac ggggcttctg	548
Leu Leu Trp Ala Gly Pro	
135 140	
agcttgctcc cctgagcctg tggctgccct ctccccagcc tggcgtggct ggggctgggg	608
gcagccttg gcccagctccg tggctgtggc ctgtggctct cactcctccc ccgacgtgaa	668
gcctccctgt ctctccgcca gctctgagtc ccaggcagct ggacatctcc aggaaaccag	728
gccatctggg caggaggcct ggggatgagg gtggggggggg acccccaggt cccggagggg	788
aagtgaagca acagcccagc tggaagggcg tcttctgcgg agaaataaag tcacttttga	848
gtcctg	854